

Evaluation of the Large Mesh Belly Panel in Small Mesh Fisheries as a Method to Reduce Yellowtail Flounder Bycatch on Southeast Georges Bank

A Report to the Northeast Cooperative Research Program

FINAL REPORT

Submitted By:

Emerson Hasbrouck – Principal Investigator
John Scotti, Tara Froehlich, Kristin Gerbino,
Joseph Costanzo, Jason Havelin, Chris Mazzeo
Cornell Cooperative Extension Marine Program
423 Griffing Avenue
Riverhead, NY 11901
(631) 727 -7850 x319

In collaboration with:

Dr. Patrick Sullivan
Cornell University
Dept. of Natural Resources
Ithaca, NY 14853
607) 255-8213

Jonathan Knight
Superior Trawl
55 State Street
Narragansett, RI 02881
(401) 782-1171

Christopher Roebuck
F/V Karen Elizabeth
Point Judith, RI

October 2014

Evaluation of the Large Mesh Belly Panel in Small Mesh Fisheries as a Method to Reduce Yellowtail Flounder Bycatch on Southeast Georges Bank

ABSTRACT

This project was developed by the Northeast Cooperative Research Program funded Squid Trawl Network to address yellowtail flounder and windowpane flounder bycatch concerns on Georges Bank by evaluating the effectiveness of a standard net modified with a large mesh belly panel to reduce the bycatch of both of these flounder species. The project was proposed by Georges Bank small mesh fishermen as means to pursue gear certification to be used for yellowtail and windowpane bycatch avoidance in Georges Bank small mesh fisheries when Accountability Measures (AM) are triggered. The Georges Bank yellowtail and windowpane flounder stocks are currently considered overfished and overfishing is occurring. The evaluation of a large mesh belly panel net in deep water while targeting squid and whiting was recommended as a bycatch avoidance solution and was conducted through this project. In response to the NEFMC's action developing Accountability Measures and sub-Annual Catch Limits for windowpane flounder as well as yellowtail flounder, quantifying windowpane bycatch reduction concurrent with yellowtail bycatch reduction was conducted through this project.

Data analysis was conducted to determine if a statistically significant difference existed in yellowtail flounder and windowpane flounder catches in the experimental net with the large mesh belly panel compared to the control net. The difference in catch of target species (squid and whiting) between the experimental net and the control net was also analyzed. Paired t-test results showed a significant difference in catch weights for yellowtail flounder and for windowpane flounder. The large mesh belly panel significantly reduced the bycatch of both flounder species. There was a 72.3% reduction in yellowtail flounder catch and 50.9% reduction in windowpane flounder catch in the net with the large mesh belly panel compared to the control net. Paired t-test results showed a non-significant result for the catch difference of whiting in the net with the large mesh belly panel compared to the control net. Paired t-test results showed a significant difference for squid. The large mesh belly net caught significantly more squid than the control net. The large mesh belly panel net retained 20% more squid than the control net. Since the experimental net did not cause significant reduction in the catch of the target species of whiting and squid but did significantly reduce bycatch of yellowtail flounder and windowpane flounder, the large mesh belly panel shows promise as a possible certified bycatch avoidance net.

INTRODUCTION

Currently, the Georges Bank (GB) yellowtail flounder stock is considered overfished and overfishing is occurring. The GB yellowtail flounder quota has been declining quite dramatically in recent years, and as a result, small-mesh discards of the stock are becoming an increasing proportion of the total U.S. catch. This project was developed to address an immediate fisheries

management need and pursue gear certification as an Accountability Measure for yellowtail bycatch in Georges Bank small mesh fisheries.

After considering the unique nature and management of the squid/whiting small mesh fishing in offshore areas, available data about relevant gear research, variability in Georges Bank yellowtail flounder catch rates on small-mesh fishery trips, the requirement to develop effective AMs in Framework Adjustment 51, and forecasts of substantially lower sub-ACLs (annual catch limits) for Georges Bank yellowtail flounder in 2014, the NEFMC Whiting and MAFMC Squid Advisory Panels made the following recommendations for management alternatives that the NEFMC should include and analyze in Framework Adjustment 51:

- Required year round use of a certified bycatch avoidance net when an AM is triggered. AM would be triggered at the end of a fishing year (April 30, 2014 at the earliest), determined a few months after the end of the fishing year, and the industry would have at least six months to procure and begin using a gear listed as an approved bycatch avoidance net at the beginning of the next fishing year (May 1, 2015 at the earliest). This timing would give industry or researchers sufficient time to evaluate experimental trawl performance. Examples of nets to be evaluated in deep water while targeting squid and whiting include a modified Ruhle trawl, a large mesh belly net, and a raised footrope trawl.

Existing research on the above nets are not directly applicable to the offshore squid/whiting fishery on Georges Bank, typically conducted using large vessels. The Ruhle trawl research was conducted using a modified squid rope trawl adapted to work with large mesh (Beutel, et al., 2008). It is not known how this net would work in the squid/whiting fishery when adapted to small mesh currently in use. The large mesh belly net has some promising features, but recent research has focused on reducing winter flounder bycatch in the inshore whiting and squid fisheries (Hasbrouck, et al., 2010, Hasbrouck, et al., 2014). Likewise, the raised footrope trawl research conducted by MADMF was completed in inshore, shallower areas and may not have the same results in deeper water with larger nets towed by larger vessels (McKiernan, et al., 1998).

As a Framework Alternative, the Council would identify a gear-based AM using approved yellowtail flounder bycatch avoidance nets that would be certified by the Regional Administrator based on submitted data and analysis of the above nets. The certification would be based on standards set by the Council in Framework Adjustment 51. If the Georges Bank yellowtail flounder AM is triggered, vessels using small-mesh trawls could only use certified yellowtail flounder bycatch avoidance nets throughout the year (NEFMC, 2013a).

Due to concerns for the declining quota, and increasing significance of small-mesh discards of GB yellowtail flounder, Framework 48 to the Northeast Multispecies Fishery Management Plan adopted a GB yellowtail flounder sub-annual catch limit (sub-ACL) for the small-mesh fisheries (NMFS, 2013). A sub-Annual Catch Limit (ACL) currently regulates small mesh fishing on

Georges Bank (GB). For the purposes of this sub-ACL, small-mesh bottom trawl fisheries are defined as those vessels that use a bottom otter trawl with a cod-end mesh size of less than 5 inches. Typical target species for vessels using this gear on GB are whiting and squid. Catches of GB yellowtail flounder by the small-mesh fisheries have generally been less than 100 mt in recent years (NEFMC, 2013b).

Recently the NEFMC council passed the following motion relative to accountability measures for small mesh fisheries on Georges Bank, to be included in Framework Adjustment 51: “To add an option as a possible Accountability Measure or as a Technical measure, any gear modifications in the small mesh fishery Georges Bank area.”

The GB yellowtail flounder quota has been declining quite dramatically in recent years and as a result, small-mesh discards of the stock are becoming an increasing proportion of the total U.S. catch. If the U.S. quota for GB yellowtail flounder is exceeded, then the U.S. quota for the following fishing year must be reduced by the amount of the overage. The pound-for-pound reduction is applied to the sub-ACL of the fishery component that caused the overage. For example, if the small-mesh fisheries caused an overage of the U.S. quota in Year 1, the small-mesh fisheries sub-ACL would be reduced by the amount of the overage in the next fishing year (Year 2). However, the small-mesh fisheries are currently required to discard all GB yellowtail flounder caught. Thus, a pound-for-pound reduction of the quota, without corresponding measures to help reduce catches of GB yellowtail flounder, would not appropriately mitigate an overage, or prevent future overages from occurring, for the small-mesh fisheries (NMFS, 2014).

Small mesh trawl nets can be modified to become highly selective in terms of the size and species of fish that they retain. Many factors influence fish capture rates including morphological and behavioral characteristics of fish as well as differences in trawl net design and construction. Successful bycatch mitigation should focus primarily on changes to the trawl design that result in applicable fishing techniques and management tools. There is an urgent need for proven methods that will work within the Georges Bank small mesh fisheries to reduce yellowtail and windowpane flounder bycatch.

The most direct option available for significant yellowtail flounder bycatch reduction in the small mesh whiting and squid fisheries is through conservation engineering and gear technological improvements. Integral to the success of any solutions that strive toward the goal of gear selectivity, is a corresponding improvement in the adoption of these methods by fishermen. This is best achieved by involving fishermen in all program aspects, from idea conception to final results. Success is also dependent on the gear modification not reducing the catch of target species (whiting and squid).

This project was developed by the Northeast Cooperative Research Program funded Squid Trawl Network (STN) to address an immediate fisheries management need and pursue gear

certification for a large mesh belly panel net to be used for bycatch reduction as an Accountability Measure for yellowtail and windowpane bycatch in Georges Bank small mesh fisheries. Discussions at the NEFMC Whiting Advisory Panel meeting in September 2013 laid the groundwork for developing gear-based AMs for Georges Bank yellowtail flounder in the small mesh fisheries. A need for proven gear concepts seeking additional consideration for small mesh trawls under this AM was the premise of this research conducted by CCE and the STN. The STN is a collaborative industry/science effort to form a comprehensive network to identify and address the challenges of bycatch and selectivity in the longfin squid fishery through innovative research. The STN was created in order to establish a collaborative industry, science and management network approach to solving the bycatch challenges of the squid fishery occurring in the Northeast. A STN Program Advisory Committee provides guidance and direction to the STN on research efforts. The STN PAC includes commercial fishing industry members, gear designers, fisheries scientists and fisheries managers. The STN PAC decided that the Squid Trawl Network would focus on an immediate response to address the yellowtail bycatch concerns on Georges Bank by evaluating the effectiveness of the large mesh belly panel on Georges Bank based on previous successful research performed by CCE in SNE/MA small mesh fisheries. Results of this previous large mesh belly panel study showed that the use of this modification resulted in an 88% reduction in winter flounder and an 83% reduction in combined demersal species (all flounders, skates, dogfish, and sea robins) (Hasbrouck, et al., 2010). These reductions were statistically significant. In addition, it should be noted that these high percentages of bycatch reduction were achieved while showing no statistically significant loss of the target species, longfin squid (Hasbrouck, et al., 2010). Similar results were proven by Milliken and DeAlteris (2004) in a project aimed at reducing flatfish bycatch in small mesh bottom trawls targeting whiting. In that project large mesh panels in the lower belly of a typical small mesh whiting net were evaluated. Their results showed large mesh belly panels proved to be effective in reducing flatfish bycatch while not reducing the catch of silver hake. Another concept considered by the STN PAC was the 12” drop chain sweep, which also showed promise in reducing winter flounder bycatch. The 12” drop chain sweep resulted in a statistically significant 78% reduction in winter flounder bycatch and a statistically significant 76% reduction in combined demersal species without a significant loss of squid (Hasbrouck, et al., 2013).

CCE maintains an excellent working relationship with fishermen from the Northeast and continually engages the commercial fishing industry, specifically the small mesh fleet, in reference to gear modifications that may be appropriate or effective in addressing bycatch of species of concern such as yellowtail and windowpane flounder. Both the 12” drop chain sweep and the large mesh belly panel modifications were designed with the collaboration of fishermen and net builders. Ultimately, it was agreed upon by the STN PAC that the large mesh belly panel modification had proven to be more effective and was to be selected for further study on Georges Bank.

It was also decided that quantifying windowpane bycatch reduction concurrent with the yellowtail bycatch reduction would be conducted. This is in response to the NEFMC's action developing Accountability Measures and sub-Annual Catch Limits for windowpane flounder as well as yellowtail flounder. Additionally, this project will extend the knowledge developed to the Georges Bank small mesh fishery and regional fisheries management councils to facilitate the transition of the application of research projects to implementation, to ensure such practices and technologies are available to managers.

Through this project, the STN aims to help resource managers and fishermen work together to sustainably use, protect, maintain and rebuild marine fisheries. More specifically, this project will develop and evaluate a conservation gear technology approach to address the issue of Georges Bank yellowtail and windowpane flounder bycatch in the small mesh fishery with the use of a large mesh belly panel net and ultimately certify this gear for approved use when AMs for small mesh fisheries are triggered. These goals will be accomplished by comparing the bycatch rates of GB yellowtail flounder and windowpane flounder for the experimental (large mesh belly panel) net and the control net as well as comparing the catch rates of the target species (whiting/squid) for each net and determining the effectiveness of the large mesh belly panel net as a successful bycatch reduction device. By definition and net design these results would also be applicable to the use of a large mesh rope trawl.


As more members of industry adopt this modification to their current trawl gear it will improve current fishing practices therefore, providing a reduction in bycatch and bycatch mortality which will allow the stocks of yellowtail and windowpane flounder to rebuild at a faster rate.

METHODS

Study Area

Two study areas have been selected for this overall project (Figure 1). The project has thus been divided into two phases to quantify gear performance in each individual area. The first is an area designated as the southern flank of Georges Bank. The second is a northern area designated as Cultivator Shoals. Observer data, NEFMC and small mesh fishermen have identified these areas as small mesh fishing areas most likely to interact with yellowtail flounder. The southern flank of Georges is a productive area fished by small mesh fishermen for squid and whiting from January – March. This is the location where the first phase experimental fishing has been completed and is the basis of this report. Due to current closures to small mesh whiting fishing on Georges Bank, the Cultivator Shoals area is closed October 31st-June 15th. Experimental fishing for phase two in this study area will occur in August 2014.

Figure 1. Map of Project Study Areas (in red) on Cultivator Shoals and the Southern Flank of Georges Bank. Green shaded areas have been closed to fishing year-round since 1994, with exceptions.



Research Design

The experimental design was intended to test the large mesh belly panel in the commercial small mesh squid and whiting fishery using existing gear and typical fishing practices. We tested for differences in both the target species catch and flounder species of concern, specifically yellowtail and windowpane flounder. We tested across appropriate identified strata of time, depth, area, and fishing practices. A single commercial twin trawl fishing vessel (F/V Karen Elizabeth) was used in this study to conduct paired replicate tows comparing a control trawl to a large mesh belly panel altered trawl (experimental trawl). This was accomplished by towing both the control and experimental nets at the same time over the same ground. A twin trawl vessel is rigged to tow two nets simultaneously in a twin-rig fashion. The study protocol used the same control/experimental trawls throughout the trip to evaluate the effectiveness of the experimental large mesh belly panel against the study objectives.

The vessel has two net reels and twin stern ramps. Both nets were set and hauled together. The vessel used one set of doors to spread the two nets (a door on each outside towing cable). The vessel used a 3-wire system with a middle winch. A “clump” (weighted sled) attached to the middle wire was towed between the two nets. Ground cables and bridles go from the clump to the inside wing of each trawl. This vessel normally tows two nets in this fashion during its

normal offshore fishing operations. Most vessels of this nature are equipped with electronic instrumentation systems that include sensors on both doors and 2 sensors on the “clump”. This allowed both nets to be fished square to the vessel, the same distance behind the vessel, and with the same wing spread. During the trip we once switched the port/starboard location of the control and experimental trawls in order to help normalize any port/starboard effect. We had an equal number of paired tows with the gear on different sides.

The control net used aboard the F/V Karen Elizabeth was an unaltered trawl net typical of the small mesh nets used in the squid and whiting fishery on Georges Bank along the southeastern area and Cultivator Shoals. The control net was a 420 x 16 cm 3-bridle 4-seam box trawl with a sweep length of 131 ft., a headrope length of 105 ft., 8" mesh (full mesh) webbing in the wings and jibs, and 6" mesh in the bunts and in the 1st bottom belly. The net had 8 cm webbing in the square, side squares, 1st top belly, 1st side bellies and the 2nd bellies. The last bellies were 6cm mesh. For the experimental trawl the 6" 1st belly was replaced with the large mesh belly. The participating captain, Captain Chris Roebuck, has extensive experience fishing for squid and whiting in the project areas.

Tow procedure had the vessel essentially fish as it would in a standard commercial fishing trip, with the exception that all tows were 30 minutes in length. The standard control net is the net that the vessel normally uses in its standard commercial squid or whiting trip. The experimental net was a standard small mesh net with the large mesh belly panel installed in the first belly. The large mesh panel is made of 80cm (32") mesh 6mm poly webbing, 2 meshes deep X 16 meshes wide sewn into the standard 16cm (6") mesh of the belly. With the ‘saw-toothing’ of the 16cm mesh, this yielded an effective opening of 3 full meshes deep, a total of about 8’ of large mesh. The panel was attached five 16cm meshes (approximately 2.5’) behind the footrope and goes from gore to gore (22 meshes wide or approximately 30’).

Number of trips and tows

This phase of the project was conducted during January 2014 in the Southern Flank of Georges Bank Study Area, near Munson Canyon. During this phase we conducted a total of 40 paired tows, all completed in one 6 day trip. All tows were 30 minutes in length and occurred during both the day and night.

On Board Catch Processing

Both nets are set and hauled together. Upon haul-back the catch from each net was kept separated on deck during the entire tow work-up procedure. The catch from each net was processed separately.

The onboard catch processing procedure followed standard NMFS survey methods as described below (NEFSC, 1988). The target was yellowtail flounder catch relative to quantifying differences in the retention between the control and experimental nets. As such, total catch of yellowtail flounder for each tow of both nets was accurately weighed. Yellowtail flounder was also sampled for length frequency. The goal was minimally 100 random length measurements per tow. When fewer individuals were caught, all were measured. We also quantified the catch of yellowtail flounder in terms of numbers as well as weights. This was accomplished by actually counting the fish (if the catch is small) or by utilizing the number of individuals in our length frequency and the weight of that sample extrapolated over the entire yellowtail flounder catch. We also quantified differences in windowpane flounder in the same manner as yellowtail flounder. Since we also wanted to quantify if the catch of whiting and squid was influenced by the experimental net modifications, the total whiting and squid catch was weighed on each tow and a length sample of at least 100 individuals was obtained. The total catch weight of all species in each tow was obtained either by direct weighing or by catch estimations. Catch estimations were based on basket or tote counts. An average weight was determined by weighing a minimum of 5 baskets or totes. Next, a count of the number of baskets or totes was made for the particular species and this number was multiplied by the average weight. This number was then recorded as the estimated total catch weight. This procedure for catch estimations, based on basket or tote counts, follows the NMFS At Sea Monitoring Program and the Observer Program Biological Sampling protocols as outlined in the NEFSC 2010 sampling manuals.

DATA ANALYSIS AND RESULTS

Below is a quantitative evaluation and summary of the data analysis. Data were analyzed primarily to determine if a significant statistical difference exists in the catch of two flounder species (yellowtail flounder and windowpane flounder) and the target species (squid and whiting) between the control and experimental nets, and to further quantify what the difference was. A variety of statistical analyses were performed on the data generated from this project. Both parametric and nonparametric statistics are used. All statistics are at the $\alpha = .05$ level.

Since only one vessel was used there was no vessel effect in the analysis relative to the catch between tows or nets. Also, since both the control and experimental nets were constructed the same (with the exception of the belly panel) and fished the same, the gear effect is only related to the belly panel installation. Statistical tests are based on pairing of the data. For each paired tow the control catch is compared to the experimental. The twin trawl design of the experiment lends itself well to pairing and a paired based analysis is the best approach. Box plots and plots of control/experimental catches by species show the distributions of each component separately (unpaired). Catch data for four key species and the catch differences between the control net and the experimental net for each tow are shown in Table 4 at the end of this report.

Unfortunately as is the case with many of these species interaction studies, it can be difficult to find commercial quantities of both target and bycatch species at the same time in the same area despite what the NMFS observer data indicates. This was the case in this study and we opted to concentrate on commercial size catches of yellowtail and windowpane flounder at the expense of smaller catches of whiting and squid.

First we looked at the difference in yellowtail flounder catch between the control net and the experimental net with the large mesh belly panel (Figures 2 and 3). Statistical analysis of the data was conducted to determine if the large mesh belly panel experimental net significantly affected retention of yellowtail flounder relative to the standard control net.

Paired t- test results showed a significant difference in the catch weight between the control and experimental net ($t = 5.7164$, $df = 39$, **p-value <0.0001**, mean of $x = 219.015$). The experimental net significantly reduced the catch of yellowtail flounder compared to the control net. The Wilcoxon test yielded similar results.

Figure 2. Boxplot distribution of yellowtail flounder catch weight in the control and experimental net






Figure 3. Distribution of Paired Tow Differences for Yellowtail Flounder



In Figure 4 below, the total weight of yellowtail flounder caught by the experimental net and by the control net for all research tows combined are compared.

Figure 4. Total Catch Weight of Yellowtail Flounder (lbs) in the Experimental and Control Net for All Trips Combined



The overall percent reduction in yellowtail flounder catch due to the large mesh belly panel treatment was 72.3% compared to the control net.

Windowpane Flounder

Next we looked at the difference in windowpane flounder catch between the control net and the experimental net with the large mesh belly panel (Figures 5 and 6). For windowpane flounder, the paired t- test results showed a significant difference in the catch weight between the control and experimental net ($t = 10.3161$, $df = 39$, **p-value <0.0001**, mean of $x = 115.32$). The experimental net caught significantly less windowpane flounder. The Wilcoxon test yielded similar results.

Figure 5. Boxplot Distribution of Windowpane Flounder Catch Weight in the Control and Experimental Net






Figure 6. Distribution of Paired Tow Differences for Windowpane Flounder



In Figure 7 below, the total weight of windowpane flounder caught by the experimental net and by the control net for all research tows combined are compared.

Figure 7. Total Catch Weight of Windowpane Flounder (lbs) in the Experimental and Control Nets for All Trips Combined



The overall percent reduction in windowpane flounder catch due to the large mesh belly panel.

Whiting

Next, the data was analyzed to determine if a significant statistical difference exists in the catch of whiting between the control and experimental nets (Figures 8 and 9). For whiting, paired t-test results showed no significant difference in the catch weight between the control and experimental net ($t = 0.1498$, $df = 39$, **p-value = 0.8817**, mean of $x = 0.4925$). The experimental net did not affect retention of whiting compared to the control net. The Wilcoxon test however did return a significant result. This is due to 1 outlier in the data. However the data are not non-normal and the outlier is relevant and part of the variability. The t-test is better in taking into account variability. Therefore the t-test is the more relevant statistic.

Figure 8. Boxplot Distribution of Whiting Catch Weight in the Control and Experimental Net






Figure 9. Distribution of Paired Tow Differences for Whiting



In Figure 10 below, the total weight of whiting caught by the experimental net and by the control net for all research tows combined are compared.

Figure 10. Total Catch Weight of Whiting (lbs) in the Experimental and Control Nets for All Trips Combined



There was no significant reduction in whiting catch due to the large mesh belly panel treatment compared to the control net.

Squid

Next, the data was analyzed to determine if a significant statistical difference exists in the catch of squid between the control and experimental nets (Figures 11 and 12). For squid, paired t- test results showed a significant difference in the catch weight between the control and experimental net ($t = -3.2734$, $df = 39$, **p-value = 0.002231**, mean of = -5.2775). The experimental net actually retained significantly more squid than the control net. The Wilcoxon test yielded similar results.

Figure 11. Boxplot Distribution of Squid Catch Weight in the Control and Experimental Net






Figure 12. Distribution of Paired Tow Differences for Squid



In Figure 13 below, the total weight of squid caught by the experimental net and by the control net for all research tows combined are compared.

Figure 13. Total Catch Weight of Squid (lbs) in the Experimental and Control Nets for All Trips Combined



Compared to the control net, the experimental net with large mesh belly panel actually retained 20% more squid. The experimental large mesh belly panel net retained on average 199 lbs. of squid per tow compared to an average of 172 lbs. retained by the control net. Although the data shows that the size of squid were on the average larger (11.56 cm) in the experimental than the control (11.33 cm), the experimental net also caught more in total number of squid and in pounds. It is difficult to speculate on why the experimental net may have retained more squid than the control net. This may be part of the randomness of the squid distribution in the ocean. Larger average size of squid in the experimental net and more squid in numbers contributes to the significant result based on pounds. We speculate that the large mesh panel had some effect on length frequency selectivity.

In summary, statistical analysis indicates that there was a significant difference in catch of both yellowtail flounder and windowpane flounder in the control net compared to the experimental net with the large mesh belly panel. The experimental net reduced the quantity of yellowtail and windowpane flounder bycatch. The overall percent reduction in yellowtail flounder catch due to the large mesh belly panel treatment was 72.3% compared to the control net. The overall percent reduction in windowpane flounder catch due to the large mesh belly panel treatment was 50.9% compared to the control net. There was no significant difference in whiting catch between the control and the experimental nets. The large mesh belly panel did not affect retention of whiting in the net. There was a significant difference in squid catch in the experimental net compared to

the control net. The large mesh belly panel net actually retained 20% more squid compared to the control net.

The catches of whiting and squid we encountered were small for commercial catches. This density level of fish is the situation that we experienced during the study. Unfortunately, as is the case in many species interaction studies, it can be difficult to find commercial quantities of both target and bycatch species at the same time in the same area despite what the NMFS observer data indicates. We opted to concentrate on commercial size catches of yellowtail and windowpane flounder at the expense of smaller catches of whiting and squid. This is part of the variability of the ocean. The findings were significantly different under the catch rates observed. In this study, sample size is the number of paired tows, not the amount of fish. The sample size we had in terms of number of coupled tows was sufficient and the experiment has enough statistical power to detect a reasonable biological difference. Larger catches may have had a different effect on those two species, but it was better to find out how the gear worked with yellowtail and windowpane flounder.

Length Frequency

Data analysis of yellowtail flounder, windowpane flounder, whiting and squid lengths was also performed to look for differences in length selectivity between the nets. The mean lengths for each tow and net were calculated for these four species. The paired differences in mean length were then compared in the control and experimental nets. Mean lengths are shown in Table 1.

Table 1. Mean Lengths (cm) of Four Species in the Control and Experimental Nets


	CONTROL	EXPERIMENTAL
Yellowtail Flounder	34.83	35.30
Windowpane Flounder	26.13	26.20
Whiting	24.65	25.17
Squid	11.24	11.57

First we conducted an ANOVA to see if there were differences in length frequency by species and different length frequencies by treatment and the effect of the species/treatment interaction term. Results of the ANOVA are in Table 2. As one would expect there are different mean lengths by species. There are also different mean lengths by treatment and, from the interaction term, the relationship in how the mean lengths differ by treatment is different by species. All results are significant. Figure 14 graphically shows the distribution of lengths by species and treatment.

Table 2. P-Value Results of ANOVA

	Mean Length
Treatment	p = <0.0001 Significant
Species	p = <0.0001 Significant
Treatment*Species	p = 0.0003 Significant

Figure 14. Boxplot of Mean Lengths by Species and Treatment




Next we conducted a series of Welch two sample t-tests. The Welch test was performed for each species to look for significant differences in length by treatment. Results are shown in Table 3 and are described below.

Table 3. Welch T-Test Results for Length Frequency Difference Between Nets

	p-value
Yellowtail Flounder	<0.0001 Significant
Windowpane Flounder	0.2990 Not Significant
Whiting	0.0016 Significant
Squid	0.0001 Significant

Figure 15 below compares the length frequency distribution for yellowtail flounder between the two nets.


Figure 15. Yellowtail Flounder Lengths as a Percent of the Total in the Control and Experimental Nets



Welch t-test results showed that yellowtail flounder were significantly larger in the experimental net ($p < 0.0001$). The average lengths for yellowtail flounder the experimental net were 0.37 cm larger than those in the control net.

Figure 16 compares the length frequency distribution for windowpane flounder between the two nets.


Figure 16. Windowpane Flounder Lengths as a Percent of the Total in the Control and Experimental Nets



Welch t-test results show that there is no significant size difference in the mean length of windowpane flounder between the control and experimental nets.


Figure 17 compares the length frequency distribution for whiting between the two nets.

Figure 17. Whiting Lengths as a Percent of the Total in the Control and Experimental Nets



Welch t-test results show a significant size difference in whiting between experimental and control nets ($p=0.0016$). Significantly larger whiting were retained in the experimental net. The mean length of whiting in the experimental net was 0.52cm larger than in the control net. Figure 18 compares the length frequency distribution for squid between the two nets.

Figure 18. Squid Length Frequency Distribution as a Percent of the Total in the Control and Experimental Nets



Welch t-test results show a significant size difference in squid between experimental and control nets ($p=0.0001$). Significantly larger squid were retained in the experimental net. The mean length of squid in the experimental net was 0.33cm larger than in the control net.


For yellowtail flounder, squid, and whiting, the size differences are significant yet they are relatively small. These statistical differences may or may not be biologically significant. However, there is a measurable difference. The fact that larger fish were retained in the net with a large mesh panel was an unexpected result.

For yellowtail flounder, not only did the experimental gear allow for significant escapement, it also seems to provide greater escapement for smaller fish. For whiting, even though there is no significant reduction in whiting catch with the experimental gear, it does seem to allow for some escapement of smaller fish. For squid, not only did the experimental gear catch more squid, they were also of larger mean length.

Day Vs. Night


Experimental fishing occurred both day and night. Although the experiment was not designed to specifically test for differences at night and differences during the day, the data was analyzed for any differences between day/night catch rates since escapement through the large mesh belly panel may have been significantly influenced by light. A Day vs. Night difference in paired tows is analyzed below (Figures 19-22).

Figure 19. Yellowtail Flounder Paired Differences for Day vs. Night




Wilcoxon rank sum test with continuity correction **p-value = 0.006119**
*significant for yellowtail flounder

Figure 20. Windowpane Flounder Paired Differences for Day vs. Night




Wilcoxon rank sum test p-value = 0.01033
*Not significant for windowpane flounder

Figure 21. Longfin Squid Paired Differences for Day vs. Night



Wilcoxon rank sum test with continuity correction p-value = 0.9432
*Not significant for squid

Figure 22. Whiting Paired Differences for Day vs. Night



Wilcoxon rank sum test with continuity correction p-value = 0.13

*Not significant for whiting

In summary, there was a statistically significant difference in the mean catch differences during the day compared to the mean catch differences at night for yellowtail flounder only. There was no significant difference for the other three species. However, this does not mean that the belly panel was not effective during the day or at night for yellowtail flounder.

Since the original analysis showed that overall for the experiment (day and night combined) the differences in the paired mean catches between the control and experimental were significant (YT, WP, SQ) and not significant (whiting), then we wanted to know if those significant and not significant results still hold up if we separate the data into 2 sub-sets: day and night. However, of the 40 total paired tows for the experiment, only 7 paired tows were at night and of those 7, two did not catch any yellowtail in either the control or experimental - so there is not enough data to say anything meaningful. However, if we look at the above plots for day/night analysis one will see that there are differences in the paired differences by day and by night for both yellowtail (significant) and windowpane (not significant). In the figures we have boxplots of each, and these are of the paired differences. Note that for both yellowtail and windowpane the differences are greater for day than they are for night. But, also note that the differences are all above zero, for both day and for night. So the overall project conclusion for yellowtail and windowpane remains the same, but it is interesting that the effect is more dramatic during the day. Even though overall escapement is significant for yellowtail and windowpane for day and night combined, the paired differences are greater during the day than at night.

DISCUSSION

For this project we looked mainly at the difference in yellowtail flounder and windowpane flounder catches in the experimental net with the large mesh belly panel compared to the control net. We also looked at the difference in catch of target species (squid and whiting) between the experimental nets and the control net. Paired t-test results showed a significant difference in catch weights for yellowtail flounder and for windowpane flounder. The large mesh belly panel significantly reduced the bycatch of both flounder species. There was a 72.3% reduction in yellowtail flounder catch and 50.9% reduction in windowpane flounder catch in the net with the large mesh belly panel compared to the control net. Paired t-test results showed a non-significant result for the catch difference of whiting in the net with the large mesh belly panel compared to the control net. Paired t-test results showed a significant difference for squid. The experimental net with the large mesh belly caught significantly more squid than the control net. The large mesh belly panel experimental net retained 20% more squid than the control net. Since the experimental net did not cause significant reduction in the catch of the target species of whiting and squid but did significantly reduce bycatch of yellowtail flounder and windowpane flounder, the large mesh belly panel shows promise as a possible certified bycatch avoidance net.

It may be true that data from twin trawl configurations sometimes does not mimic single trawl configurations but it does offer a direct side-by-side comparison of different gear types. Further work using single trawls using replicate tows or side-by-side vessels could add additional data to strengthen already proven results.

SUMMARY OF CONCLUSIONS

- The large mesh belly panel has proven to be functionally effective in significantly reducing the quantity of yellowtail flounder bycatch. The large mesh belly panel reduced yellowtail flounder bycatch by 72.3%.
- There was no significant difference in whiting catch between the control net and the net modified with the large mesh belly panel. Retention of this target species was maintained using the experimental net.
- The large mesh belly panel has also proven to be functionally effective in significantly reducing the quantity of windowpane flounder bycatch. The large mesh belly panel reduced windowpane flounder bycatch by 50.9%.
- There was a significant difference in squid catch between the control net and the net modified with the large mesh belly panel. The experimental net retained 20% more squid compared to the control net.

Table 4. Catch Data for Four Key Species for All Tows and Difference Between Control and Experimental Nets

TRIP NUMBER	TOW NUMBER	CONTROL NET CATCH (LBS)				EXPERIMENTAL NET CATCH (LBS)				DIFFERENCES (CONTROL – EXPERIMENTAL)			
		YELLOWTAIL	WINDOWPANE	SQUID	WHITING	YELLOWTAIL	WINDOWPANE	SQUID	WHITING	YELLOWTAIL DIFFERENCE	WINDOWPANE DIFFERENCE	SQUID DIFFERENCE	WHITING DIFFERENCE
1	1	0	27.5	36.6	3.4	0	7.9	56.4	1.8	0	19.6	-19.8	1.6
1	2	3.8	31.6	81	4.5	1.3	15.2	61.7	1.7	2.5	16.4	19.3	2.8
1	3	5.2	53.5	182.9	28.9	3.2	30.3	202.9	23.4	2	23.2	-20	5.5
1	4	0	6.2	93.2	55.6	0.8	3.2	119.4	174	-0.8	3	-26.2	-118.4
1	5	0	0	25.5	256.6	0	0.8	32.9	222.8	0	-0.8	-7.4	33.8
1	6	0	0.5	75.8	51.7	0	0	103.1	81.6	0	0.5	-27.3	-29.9
1	7	44.9	180.7	18	16.3	11.6	97.2	23.3	13.1	33.3	83.5	-5.3	3.2
1	8	99.5	452	26.6	20.7	21.4	255.8	38.7	8.7	78.1	196.2	-12.1	12
1	9	223.4	281	20.7	6.3	46.1	56.1	24.4	5.5	177.3	224.9	-3.7	0.8
1	10	329.4	381.8	30.9	8.5	82.5	177.3	22.7	4.9	246.9	204.5	8.2	3.6
1	11	419.8	246.1	13.7	23.4	123.7	134.4	20.3	13.7	296.1	111.7	-6.6	9.7
1	12	204.7	159.4	9.4	12.9	74	83.9	25.3	6.2	130.7	75.5	-15.9	6.7
1	13	311.3	289.6	15.7	7.2	70.1	134.1	32.5	6.9	241.2	155.5	-16.8	0.3
1	14	336.9	311.5	18.6	16	47.6	123.3	15.8	6.3	289.3	188.2	2.8	9.7
1	15	311.7	439.2	20	13.2	90.3	258.1	30	5.6	221.4	181.1	-10	7.6
1	16	246.2	63.7	10.5	4.8	157.4	77.9	13.7	5.1	88.8	-14.2	-3.2	-0.3
1	17	326.2	126.8	6.5	4.9	168	78.1	9.2	4.1	158.2	48.7	-2.7	0.8
1	18	91	215.3	6.4	9.7	140.7	149.4	12.1	8.9	-49.7	65.9	-5.7	0.8
1	19	305.1	346.1	12.4	8	104.9	194.7	14.2	3.7	200.2	151.4	-1.8	4.3
1	20	246.5	255.6	13.4	5.1	65.4	142.8	14.1	2.2	181.1	112.8	-0.7	2.9
1	21	243.7	313.8	14.8	7.3	27.6	71.3	44.6	8.5	216.1	242.5	-29.8	-1.2
1	22	245.8	233.6	34.7	6.9	56.2	128.6	32	2.6	189.6	105	2.7	4.3
1	23	278.4	196.4	11.6	15.8	51.1	70.8	18.6	10.4	227.3	125.6	-7	5.4
1	24	343.9	160.4	17.9	9.5	87.6	94.8	12.4	6.5	256.3	65.6	5.5	3
1	25	170.1	279.5	12.5	6.2	54.8	120.7	17.4	2.4	115.3	158.8	-4.9	3.8
1	26	345.8	356.5	9.6	10.1	55.7	82.4	18.1	5.2	290.1	274.1	-8.5	4.9
1	27	149.7	266.9	20.9	4.1	64.2	158.9	20.9	5.5	85.5	108	0	-1.4
1	28	324.5	349.8	14.1	12.3	95.8	144.6	26.8	6.3	228.7	205.2	-12.7	6
1	29	223.4	295	13.5	8.7	91.2	199	14.3	4.4	132.2	96	-0.8	4.3
1	30	356.3	319.3	9.6	10.8	176.9	150.7	7.3	7	179.4	168.6	2.3	3.8
1	31	164.8	354.1	9.4	7.6	77.2	246.9	7	6.2	87.6	107.2	2.4	1.4
1	32	302.5	405.1	6.9	7.3	118.7	274	8.8	3.5	183.8	131.1	-1.9	3.8
1	33	274.5	169.9	14.5	5.1	35.7	52.1	14.6	2.1	238.8	117.8	-0.1	3

1	34	370.8	212.3	15.4	3.5	64.8	92.8	13.5	3.5	306	119.5	1.9	0
1	35	300.5	185.7	12	4.5	44	56.1	18.9	2.3	256.5	129.6	-6.9	2.2
1	36	503.9	227.5	12.5	5.1	97.3	96.5	19.2	2.1	406.6	131	-6.7	3
1	37	387	251.8	18.9	5.4	72.3	119	24.5	4	314.7	132.8	-5.6	1.4
1	38	706.7	238.4	16.7	7.4	75	83.4	18.9	5.1	631.7	155	-2.2	2.3
1	39	1135.9	210.1	14.6	8.8	336.4	81.3	9.1	4.1	799.5	128.8	5.5	4.7
1	40	1775.3	160.4	32.8	10.6	457	97.4	22.2	3.1	1318.3	63	10.6	7.5

LITERATURE CITED

- Beutel, D., L. Skrobe, K. Castro, P. Ruhle Sr., P. Ruhle Jr., J. O'Grady, J. Knight. 2008. Bycatch reduction in the Northeast USA directed haddock bottom trawl fishery. *Fisheries Research* 94.2 (2008): 190-198.
- Hasbrouck, E., P. Sullivan, J. Knight, C. Weimar, and B. Brady. 2012. A Method To Reduce Winter Flounder Retention Through The Use Of An Avoidance Gear Adaptation In The Small Mesh Trawl Fishery Within The Southern New-England/Mid-Atlantic Winter Flounder Stock Area. Final Report to Commercial Fisheries Research Foundation.
- Hasbrouck, E., J. Knight, R. Jones, P. Ruhle Jr., K. Ketcham, P. Sullivan, W. Grimm, and S. Arnold. 2013. An Evaluation Of The Avoidance Gear 12" Drop Chain Sweep As A Method To Reduce Winter Flounder Retention In The Small Mesh Squid Trawl Fishery Within The SNE/MA Winter Flounder Stock Area. Final Report to Commercial Fisheries Research Foundation.
- Hasbrouck, E., J. Knight, P. Sullivan, P. Merris and J. Wise. 2014. Gear Trials – Reduction of Winter Flounder Bycatch: Proof of Concept Program – Experimental Fishery Phase. Final Report to the Commercial Fisheries Research Foundation.
- McKiernan, D.J. R. Johnston, B. Hoffman, H. A. Carr, H.O. Milliken and D. McCarron. 1998. 1998 Southern Gulf of Maine Raised Footrope Trawl in the Whiting Fishery. *Comm. Of Mass Div. Mar. Fish.* 30 pp.
- Milliken, H.O., J.T. DeAlteris. 2004. Evaluation of a Large-Mesh Panel to Reduce the Flatfish Bycatch in the Small-Mesh Bottom Trawls Used in the New England Silver Hake Fishery. *North American Journal of Fisheries Management*, Volume 24, pp 20-32.
- NMFS (National Marine Fisheries Service). 2013. Magnuson-Stevens Fishery Conservation and Management Act Provisions; Fisheries of the Northeastern United States; Northeast (NE) Multispecies Fishery; Framework Adjustment 48. *Federal Register / Vol. 78, No. 86*
- NMFS (National Marine Fisheries Service). 2014. Magnuson-Stevens Fishery Conservation and Management Act Provisions; Fisheries of the Northeastern United States; Northeast Groundfish Fishery; Framework Adjustment 51. *Federal Register/ Vol. 79, No. 77*
- NEFMC. 2013a. Memorandum From NEFMC Whiting and MAFMC Squid Advisory Panels (Aps) Subject: Accountability measure alternatives in Northeast Multispecies Framework 51. September 9, 2013.
- NEFMC. 2013b. Memorandum From: Andrew Applegate To: Whiting and Squid Advisors. Subject: Background Information About Georges Bank Yellowtail Flounder Accountability Measures. September 3, 2013.
- NEFSC. 1988. An evaluation of the bottom trawl survey program of the Northeast Fisheries Science Center. NOAA Tech. Mem. NMFS/NEC-52, 82 pp.

APPENDIX I

Review of "Evaluation of the Large Mesh Belly Panel in Small Mesh Fisheries as a Method to Reduce Yellowtail Flounder Bycatch on Southeast Georges Bank"

Reviewer Comments and Response to Reviewer's Comments

Note - It should be noted that this draft report is part of a larger project to evaluate the potential bycatch reduction of the large mesh belly panel in small mesh fisheries. Sampling in other areas is continuing and will be analyzed in additional reports when completed.

The final report has been updated to include the information in this review.

Reviewer 1

Introduction

The report outlines the rationale for the study and the funding in great detail and the importance of the outcome is clear to the reader. Quite a bit of previous work on the Ruhle trawl, the large mesh belly net, and raised footrope trawl is briefly discussed and seems quite relevant, however references allowing the reader to examine this previous work would be very helpful here. Although it is clear that the gear design was worked out cooperatively with the fishing industry, a more thorough discussion of the previous work, what alternatives were considered and how the gear design used was arrived at would be very useful here. Why was this particular design chosen over the various alternatives?

Response:

Discussions at the NEFMC Whiting Advisory Panel meeting in September 2013 laid the groundwork for developing gear-based AMs for Georges Bank yellowtail flounder in the small mesh fisheries. A need for proven gear concepts seeking additional consideration for small mesh trawls under this AM was the premise of this research conducted by CCE and the STN. Discussions with the STN Program Advisory Committee (PAC) directed the research conducted in this study. The STN PAC includes commercial fishing industry members, gear designers, fisheries scientists and fisheries managers. The STN PAC decided to evaluate yellowtail flounder bycatch reduction with the large mesh belly panel on Georges Bank based on previous successful research performed by CCE in SNE/MA small mesh fisheries. Results of this previous large mesh belly panel study showed that the use of this modification resulted in an 88% reduction in winter flounder and an 83% reduction in combined demersal species (all flounders, skates, dogfish, and sea robins) (Hasbrouck, et al., 2010). These reductions were statistically significant. In addition, it should be noted that these high percentages of bycatch reduction were achieved while showing no statistically significant loss of the target species, longfin squid (Hasbrouck, et al., 2010). Similar results were proven by Milliken and DeAlteris (2004) in a project aimed at reducing flatfish bycatch in small mesh bottom trawls targeting whiting. In that project large mesh panels in the lower belly of a typical small mesh whiting net were evaluated. Their results showed large mesh belly panels proved to be effective in reducing flatfish bycatch while not reducing the catch of silver hake.

Another concept considered by the STN PAC was the 12" drop chain sweep, which also showed promise in reducing winter flounder bycatch. The 12" drop chain resulted in a statistically significant 78% reduction in winter flounder bycatch and a statistically significant 76% reduction in combined demersal species without a significant loss of squid (Hasbrouck, et al., 2013). Ultimately, it was agreed upon by the STN PAC that the large mesh belly panel modification had proven to be more effective and was to be selected for further study on Georges Bank.

Literature Cited:

- Beutel, D., L. Skrobe, K. Castro, P. Ruhle Sr., P. Ruhle Jr., J. O'Grady, J. Knight. 2008 "Bycatch reduction in the Northeast USA directed haddock bottom trawl fishery." *Fisheries Research* 94.2 (2008): 190-198.
- Hasbrouck, E., P. Sullivan, J. Knight, C. Weimar, and B. Brady. 2012. A Method To Reduce Winter Flounder Retention Through The Use Of An Avoidance Gear Adaptation In The Small Mesh Trawl Fishery Within The Southern New-England/Mid-Atlantic Winter Flounder Stock Area. Final Report to Commercial Fisheries Research Foundation.
- Hasbrouck, E., J. Knight, R. Jones, P. Ruhle Jr., K. Ketcham, P. Sullivan, W. Grimm, and S. Arnold. 2013. An Evaluation Of The Avoidance Gear 12" Drop Chain Sweep As A Method To Reduce Winter Flounder Retention In The Small Mesh Squid Trawl Fishery Within The SNE/MA Winter Flounder Stock Area. Final Report Summary.
- Hasbrouck, E., J. Knight, P. Sullivan, P. Merris and J. Wise. 2014. Gear Trials – Reduction of Winter Flounder Bycatch: Proof of Concept Program – Experimental Fishery Phase. Final project report to the Commercial Fisheries Research Foundation.
- McKiernan, D.J. R. Johnston, B. Hoffman, H. A. Carr, H.O. Milliken and D. McCarron. 1998. 1998 Southern Gulf of Maine Raised Footrope Trawl in the Whiting Fishery. *Comm. Of Mass Div. Mar. Fish.* 30 pp.
- Milliken, H.O., J.T. DeAlteris. 2004. Evaluation of a Large-Mesh Panel to Reduce the Flatfish Bycatch in the Small-Mesh Bottom Trawls Used in the New England Silver Hake Fishery. *North American Journal of Fisheries Management*, Volume 24, pp 20-32

The above discussion will be added to the Introduction section of the final report and the cited references will be added to the Literature Cited section.

Study Design

The project was well-conceived with a very solid design to test the effects of the large mesh belly panel. The twin trawl allowed the elimination of almost all extraneous variables and allowed very high power in testing the difference between control and treatment. The number of replicate tows seems to have been sufficient given the large differences between control and treatment and should be improved by the continued work in August of 2014. Although there does not appear to have been any difference between port and starboard (and there is no reason to believe that this would be the case given the study design), it would have been better to alternate sides more often to sufficiently test this hypothesis.

Data analysis

The tests used to analyze the data are appropriate for the methods and were quite effective in elucidating the differences in catches with the large mesh panel. Although I don't think it would make much difference in this case, it would have been nice to see a test comparing the actual length frequencies between treatment and control for each of the species (K-S or Cramer-von Mises test or perhaps a simple modeling effort) in addition to the comparison of mean lengths via the Welch t-tests. There could be cases where these tests would tell a very different story, although looking at the length-frequency data, I don't believe that to be the case here, with the possible exception of the whiting.

Response:

The K-S test and the Cramer-von Mises test are used to compare the distributions of two groups by looking at differences in the empirical cumulative distributions between groups. These tests are most helpful in situations where the means of the distributions look similar, but the shape of the distribution of one or the other group looks different (e.g. one appears skewed, while the other does not, or one appears "lumpy"). In these situations the standard tests applied to means or even variances might not work so well in detecting differences. In general, though, using the

mean to check to see if the distributions are different is the first step to take and is often the simplest one to employ and interpret. In our case, using these simple tests we found that 3 of the 4 species examined showed a statistically significant difference in the mean. We think this is sufficient to demonstrate the point that the length distributions are different between treatment and control.

Discussion

I would like to see some discussion as to why the experimental net caught more squid. I think this is a very interesting finding and deserves some speculation as to why this might be. Is the detected size difference alone enough to account for this difference? It is also not clear to me why the experimental gear would retain larger squid on average. The differences in the whiting catches also deserve some discussion. Although the overall catch weights were not significantly different there appears to be a clear difference in the size selectivity between the two treatments.

Response:

The experimental large mesh belly panel net retained on average 199 lbs. of squid compared to an average of 172 lbs. retained by the control net. Although the data shows that the size of squid were on the average larger (11.56 cm) in the experimental than the control (11.33 cm), the experimental also caught more in total number of squid and in pounds. It is difficult to speculate on why the experimental net may have retained more squid than the control. This may be part of the randomness of the squid distribution in the ocean. Larger average size of squid in the experimental and more squid in numbers contributes to the significant result based on pounds. We speculate that the large mesh panel had some effect on length frequency selectivity. However this experiment was not designed as a size selectivity study.

We will add this discussion to the report.

I also think some discussion of what other gear modification possibilities might be possible and why the authors believe the method chosen is superior to the other potential methods would be very helpful here. It is perhaps beyond the scope of this study, but some discussion of how the use of this gear would affect yellowtail and windowpane populations of Georges Bank relative to current practices would be useful for the reader.

Response:

CCE maintains an excellent working relationship with fishermen from the Northeast and continually engages the commercial fishing industry, specifically the small mesh fleet, in reference to gear modifications that may be appropriate or effective in addressing bycatch of species of concern such as yellowtail and windowpane flounder. Both the 12" drop chain sweep and the large mesh belly panel modifications were designed with the collaboration of fishermen and net builders. These are ideas that they may have tried with on their own in some variation. CCE took all these ideas and narrowed it down to a single design that is easily modified to existing gear, is effective in reducing flounder bycatch and is easily compliant to management and enforcement standards. Yes, it may be beyond the scope of this study but one of the main goals of conservation gear technology is to help rebuild stocks in decline by reducing bycatch of those species. The large mesh belly panel has proven to be effective in reducing bycatch of windowpane and yellowtail flounder on Georges Bank. As more members of industry adopt this modification to their current trawl gear it will improve current fishing practices therefore, providing a reduction in bycatch and bycatch mortality which will aid in the stocks of yellowtail and windowpane flounder to rebuild faster. See also the above response to Reviewer 1 first comment.

Overall this study was well-conceived and executed. They showed a very significant decrease in the catches of yellowtail and windowpane flounder and the results seem very promising in terms of meeting the goals of reducing bycatch of these species in the small mesh fisheries, while not substantially affecting the catch rates and profitability of the participating vessels.

Reviewer 2

This research is certainly worthwhile and there is valid need for a modified gear to reduce yellowtail and windowpane flounder catches in the squid and whiting small mesh fishery on Georges Bank. The results contained in the report show that the large mesh belly panel was effective at reducing catch of the bycatch species, YT and WP flounders, and should be considered a potential option as a bycatch avoidance net. However, I am not convinced that the study encountered enough of the target species, squid and whiting, to definitively state no statistical difference in retention of the target species between experimental and control nets. The study only caught approximately 1400lbs of whiting and 2200lbs of squid, compared to 15000lbs yellowtail and 13000lbs windowpane.

Response:

The findings were significantly different under the catch rates observed. Whether they are different under other catch rates is not something we can show. This is not a matter of sample size. The sample size we had in terms of number of coupled tows was sufficient. The challenge really is that if the density of fish in the water is greater (or even less) will there be some kind of density dependent effect that obscures the differences we saw in this experiment. But, this density level of fish is the situation that we experienced during the study. Unfortunately as is the case in many of these species interaction studies it can be difficult to find commercial quantities of both target and bycatch species at the same time in the same area despite what the NMFS observer data indicates. This was the case here and we opted to concentrate on commercial size catches of yellowtail and windowpane flounder at the expense of smaller catches of whiting and squid. This is part of the variability of the ocean. So, yes the catches of whiting and squid are smaller than commercial catches. Larger catches may have had a different effect on those two species, but it was better to find out how the gear worked with yellowtail and windowpane flounder. We will add an explanation in the report about the small catches of whiting and squid.

One other area of concern regarding the results is the potential differences in area swept between the experimental and control gears. The report states that net mensuration data was collected, however, these data are not compared between gears in the report. There is significant potential for spread differences, and therefore area swept differences, between the control and modified nets that could contribute to differences in catch rates. Net mensuration data may also help to explain the differences in catch between port and starboard sides observed in yellowtail.

Response:


We did not analyze the net mensuration data. However, when taking a simple quantitative look at the data, there is no difference in door spread between the experimental and control nets for most of the 39 tow pairs. Door spread was normally 37 fathoms. In the instances where door spread varies between experimental and control tows, it does not vary by more than one fathom. However, if we modify a net and it catches what it is supposed to and doesn't catch what it is not supposed to, does the mensuration matter? Perhaps adding the panel causes it to open more or less or differently. That information might be good to know from the perspective of understanding the physics of how the net is operating. However, it doesn't affect the conclusions from the statistical tests. This project was focused on conservation gear technology and

producing an effective gear modification to reduce yellowtail and windowpane flounder bycatch, this study was not designed to address swept area biomass.

I have concerns regarding any differences between day and night catch rates. The report states that fishing occurred both day and night and escapement through the large mesh belly panel may be significantly influenced by light. The data should be analyzed for any differences between day/night catch rates. If this gear is ultimately going to be considered for use as bycatch avoidance gear, the escapement/catch rates need to be understood for both day and night fishing.


Response:

Day Vs. Night is analyzed below.




Wilcoxon rank sum test with continuity correction
p-value = 0.006119 *significant for yellowtail flounder

Yellowtail flounder is the only one that is significant.




Wilcoxon rank sum test

p-value = 0.01033 *Not significant for windowpane flounder



Wilcoxon rank sum test with continuity correction

p-value = 0.9432 *Not significant for squid



Wilcoxon rank sum test with continuity correction
 p-value = 0.13 *Not significant for whiting

In summary, there was a statistically significant difference in the mean catch differences during the day compared to the mean catch differences at night for yellowtail only. There was no significant difference for the other three species. However, this does not mean that the belly panel was not effective during the day or at night for yellowtail flounder.

Regardless of whether there is a difference in the mean catch differences during the day compared to the mean catch differences at night, the more appropriate question may be: Since the original analysis showed that overall for the experiment (day and night combined) the differences in the paired mean catches between the control and experimental were significant (YT, WP, SQ) and not significant (whiting), then do those significant and not significant results still hold up if we separate the data into 2 sub-sets: day and night. However, of the 40 total paired tows for the experiment, only 7 paired tows were at night and of those 7, two did not catch any yellowtail in either the control or experimental - so there is not enough data to say anything meaningful. The experiment was not designed to specifically test for differences at night and differences during the day. However, if we look at the above plots for day/night analysis one will see that there are differences in the paired differences by day and by night for both yellowtail (significant) and windowpane (not significant). In the figures we have boxplots of each, and these are of the paired differences. Note that for both yellowtail and windowpane the differences are greater for day than they are for night. But, also note that the differences are all above zero, for both day and for night. So the overall project conclusion for yellowtail and windowpane remains the same, but it is interesting that the effect is more dramatic during the day. We will note in the report that for yellowtail and windowpane that even though overall escapement is significant for day and night combined, the paired differences are greater during the day than at night.

Reviewer 3

In my opinion, the report reads well and describes the work that was completed. I have a few comments below on the results and discussion section of the report. As I do not have a strong knowledge of the management systems/plans, I did not review the introduction section of this report.

Comments/suggestions:

Major:

1. I suggest that the report provide a better explanation of the analysis of the side differences in Tables 1,2,3,4 and Figures 5,9,13,17. From the methods section, the nets were switched once halfway through the study. Was the difference an analysis based on the mean of the differences between the nets when they were on one side versus the other? I expect that the analysis is an examination of the catch rates between the nets when they were on one side of the vessel compared to when they were on the alternate side. In either case, the analysis would be subject to the influence of sampling different populations of fish and is likely meaningless as a method to compare side effects. I am not sure how you would measure a side effect using the experimental design outlined in this report. Regardless, the methods should be outlined and the researchers need to consider if it is worth reporting if it might lead the reader to a misinformed conclusion.

Response:

The original study design was to switch the side of the experimental and control nets at the end of each day in order to randomize for side. However, due to extenuating circumstances while fishing related to complications involved with switching nets, the nets were switched for side only once during the experiment. However, we have an approximately equal number of tows with the experimental net on the port side as on the starboard side. Therefore it became important for us to at least explore the possibility that side made a difference. In the end, the non-significant findings for three species indicate that side seemed to play a minor role in influencing our findings. We agree with the reviewer that including the side analysis might lead the reader to confusion about the issue and a misinformed conclusion. Therefore we have removed the analysis and discussion of side from the report.

2. The catch rates for the whiting and squid appear to average around 10 lbs /tow. These are not commercial quantities of catch. I understand that the researchers were interested in determining the loss of yellowtail and windowpane flounder but I suggest that the authors comment that the targeted catch rates were low and caution about conclusions based on these numbers. In fact, the low sample numbers are likely the reason for the 20% higher catch rate of squid in the net with the large mesh belly panel. If the gear is the same, i.e., same net opening, there is no reason that the experimental net should show a significant increase in catch. Additionally, a power analysis could be performed to determine the adequacy of the data. The authors could have referenced other work on this panel design to support the conclusion about targeted catch retention, but without additional information, I am reticent to believe the data on the targeted catch rates. The discussion should frame this so that the reader is informed about the results of this work.


Response:

In this study, sample size is the number of paired tows, not the amount of fish. The experiment has enough statistical power to detect a reasonable biological difference. Examining how the density of fish in the water plays a role cannot be explored via a power analysis. That would be a nonlinear density dependent effect unrelated to the sample sizes we have been using in this experiment. In other words, our sample size is appropriate, we would just need to apply the experiment to different densities of different components of the catch. This isn't really a design problem so much as exploring other factors that may influence our findings. That would be a

series of different experiments since lots of external factors can affect our findings. This doesn't affect the quality of our statistical design. If one wants to know what will happen at other fish densities, then we will need to have another experiment. But, then we might wish to have, for example, high:high density, high:low density, low:high density and low:low density in terms of the fish we wish to avoid relative to those we wish to catch to cover all the bases. The comment about fish density is a decent one, but we cannot test against all possibilities of density of catch. We acknowledge that we cannot control for all variability in the ocean. However, we attempted to randomize the variability and control the things we can control. See also our response to Reviewer 2 above on a closely related comment.


Minor:

4. In my opinion, the information on Figures 3,7,11,15 could be better portrayed with a scatter plot of the catch of the experimental net plotted against the catch of the control net with a line through the axis at 50% (example below). This would provide the dispersion and frequency of the differences.




Response:


This is an interesting way to present the data. One thing that is captured in these plots is the actual magnitude of the catches by each net. But, it is difficult in this plot to gauge the actual paired differences between the points, which is what the test is based on and what the histograms provided in the report show. However, if one looked at the perpendicular distance from the one-to-one line to the point, then this distance would be $\sqrt{2 * (\text{catch1} - \text{catch2})^2}$ or $(\sqrt{2}) * \text{catch difference}$, so it would be at least proportional to the catch difference that we are measuring in the test, but we think the histograms do a better job of this. Scatterplots suggested by the reviewer are shown below. However, we believe the histograms in the report do a better job of showing the paired differences in catch, which is what the statistical tests are based on. We think it is more appropriate to retain the plots we have in the report.




The above figure corresponds to Figure 3 on Yellowtail.



The above figure corresponds to Figure 7 for Windowpane.



The above figure corresponds to Figure 11 for Whiting.



The above figure corresponds to Figure 15 for Squid.

5. Bar for large mesh belly panel in Figure 4 is narrow.

Response:

The figure looks normal in our version of the report.

6. No previous studies and comparison to previous studies are reported in this report.

Response:

See response to Reviewer 1.

7. Data from twin trawl configurations sometimes do not mimic single trawl configurations.

The discussion should mention this and suggest that further work on vessels towing single trawls would help verify these results.

Response:

It can be true that data from twin trawl configurations sometimes does not mimic single trawl configurations but it does offer a direct side-by-side comparison of different gear types. Further work using single trawls using replicate tows or side-by-side vessels could add additional data to strengthen already proven results.

8. A table with the catch data would help in understanding the results. As there were only 40 hauls, the data could be provided as a table instead of an `x` appendix.

Response:

The table is now part of the report.