

Juveniles and adults were distinguished based on lengths-at-maturity for each species, which was defined according to the length at which 50% of the fish in a population mature sexually. For most species, these sizes vary by sex and stock units. They also vary over time, according to changes in growth rate, sometimes considerably. Lengths used to distinguish juveniles and adults for most species were based on data reported by O'Brien et al. (1993). Lengths at maturity for the skate species were based on information included in EFH source documents. These lengths are listed in Table 1. In most cases, O'Brien et al. based 50% lengths at maturity on females; if there was more than one size available because of analyses that were performed at different time periods or for different stocks, they were averaged.

$$r(l) = \{ \exp(a + bl) / [1+\exp(a + bl)] \}$$

Table 1 – Lengths-at-maturity used to distinguish juveniles and adults in EFH designations. Juveniles are less than the specified length; adults are equal to or larger.

<i>Species</i>	<i>Length (cm) at 50% Maturity O'Brien et al. (1993) and EFH Skate Source Document</i>	<i>Length (cm) at maturity (rounded to nearest 5 cm for analysis of juvenile and spawning distributions)</i>		<i>Approximate length (rounded up to 5 cm increment) at greater than 80% Maturity from 2002-2012 spring and fall trawl survey data</i>		
		<i>Calculated from parameters in latest assessment, generally GARM III</i>	<i>Red values are average L20/L50 and L80/L50 ratios of other species</i>	<i>L20</i>	<i>L50</i>	<i>L80</i>
American Plaice	27	23.6 (25)		27.6	31.6 (30)	30
Atlantic Cod	35	35.4-36.8 (35)		43-44.5	49.2-53.6 (50)	50
Atlantic Herring	25	(20)		NA	(25)	25
Barndoor Skate	102	(85)		NA	(115)	115*
Clearnose Skate	61	(50)		NA	(70)	
Deep-sea Red Crab	8			NA		
Goosefish	43	(35)		NA	(45)	45
Haddock	32	28.2-28.3 (30)		33-34.7	37.8-41.1 (40)	40
Little Skate	50	(45)			(55)	
Ocean Pout	29					
Offshore Hake	30	(25)			(35)	
Pollock	39	38.8 (40)		45.4	51.9 (50)	45
Red Hake	26	(20)			(35)	35
Redfish	22	19.2 (20)		22.0	24.8 (25)	25

Species	Length (cm) at 50% Maturity O'Brien et al. (1993) and EFH Skate Source Document	Length (cm) at maturity (rounded to nearest 5 cm for analysis of juvenile and spawning distributions)		Approximate length (rounded up to 5 cm increment) at greater than 80% Maturity from 2002-2012 spring and fall trawl survey data
		Calculated from parameters in latest assessment, generally GARM III	Red values are average L20/L50 and L80/L50 ratios of other species	
		L20	L50	L80
Rosette Skate	46	(40)		(55)
Sea Scallop	10			
Silver Hake	23	(20)		(30) 30
Smooth Skate	56	(50)		(65)
Thorny Skate	84	(70)		(95)
White Hake	35	25.0 (25)	35.1	45.2 (45) 60
Windowpane	22	17.5-18.2 (20)	20.5-21.3	23.5-24.4 (25)
Winter Flounder	27	26.7 (25)	29-29.1	31.1 (30) 30
Winter Skate	85	(70)		(95)
Witch Flounder	30	28.1 (30)	32.9	31.1 (40) 40
Yellowtail Flounder	27	24.6-25.8 (25)	27.4-28.2	30.2-30.7 (30) 30

Wolfish – 47 cm (Templeman 1986)

Table 2. Cumulative proportion of abundance at age by species, survey, and stock area. First line of data represents an approximate L20 for each species. Second line of data represents a size that approximates the 90th percentile of age 1 fish (some species use age 2) for the predominate stock area for each species.

Species	Survey	Length (cm)	Spring 2002-2012		Region		Fall 2002-2011			Age		Mid-Atlantic			Georges Bank			Gulf of Maine			Scotian Shelf					
			0			1			2			3			0			1			2					
American plaice	Spring	25													100.0%	99.4%	63.4%	100.0%	99.5%	86.3%	100.0%	98.6%	85.1%			
		12													91.5%	4.3%	0.0%	84.2%	2.6%	0.0%	90.5%	0.0%	0.0%			
	Fall	25													100.0%	86.9%	37.7%	100.0%	98.1%	65.7%	100.0%	100.0%	97.1%	57.1%		
		18													100.0%	89.6%	16.4%	1.0%	100.0%	98.2%	35.4%	1.7%	100.0%	98.0%	16.2%	0.0%
Atlantic cod	Spring	35	100.0%	100.0%	15.0%	0.0%	100.0%	99.7%	29.5%	0.8%	100.0%	100.0%	75.9%	12.1%	100.0%	100.0%	45.8%	2.2%								
		24	100.0%	41.4%	0.0%	0.0%	100.0%	65.2%	0.7%	0.0%	100.0%	90.6%	14.7%	0.0%	100.0%	95.3%	1.0%	0.0%								
	Fall	35	100.0%		0.0%	0.0%	100.0%	66.4%	2.0%	0.0%	100.0%	94.0%	29.2%	2.7%	100.0%	84.9%	13.3%	2.7%								
		34	100.0%		0.0%	0.0%	100.0%	58.6%	1.4%	0.0%	100.0%	91.3%	25.3%	1.5%	100.0%	80.2%	10.7%	0.0%								
Atlantic herring	Spring	20	100.0%	99.5%	65.5%		100.0%	99.8%	73.1%	100.0%	100.0%	99.8%	75.8%		100.0%	100.0%	70.6%									
		9	100.0%	0.3%	0.0%			91.7%	0.2%	0.0%	100.0%	94.1%	1.1%	0.0%			90.0%	7.1%	0.0%							
	Fall	20	100.0%	84.2%	66.7%		100.0%	81.8%	12.0%		100.0%	84.1%	11.8%			100.0%	90.6%	10.2%								
		16	100.0%	0.0%	0.0%			95.8%	3.2%	0.0%		96.9%	8.1%	0.0%			100.0%	10.4%	0.0%							
Goosefish	Spring	35		100.0%	100.0%				100.0%	100.0%		100.0%	100.0%			100.0%				100.0%						
		28		100.0%	84.2%				100.0%	92.3%		100.0%	100.0%			93.0%				100.0%						
	Fall	35	100.0%	100.0%	100.0%		76.9%	100.0%	100.0%		54.5%	100.0%	100.0%		70.0%	100.0%	100.0%		100.0%							
		26	100.0%	100.0%	90.9%	0.0%	100.0%	100.0%	95.2%	0.0%	100.0%	100.0%	77.8%	0.0%	100.0%	100.0%	100.0%		100.0%							
Haddock	Spring	30	100.0%	0.0%				99.9%	48.0%	7.5%		100.0%	35.9%	3.5%		100.0%	56.9%	11.7%								
		24	67.4%	0.0%				88.6%	7.8%	0.0%		93.3%	1.4%	0.0%			95.0%	6.4%	3.3%							
	Fall	30	100.0%	0.0%			100.0%	68.5%	10.5%	0.2%	100.0%	77.4%	4.9%	0.4%	100.0%	83.9%	8.1%	0.0%								
		34	100.0%	100.0%			100.0%	93.5%	29.3%	7.8%	100.0%	97.4%	27.6%	3.3%	100.0%	99.1%	45.1%	9.4%								
Ocean pout (all years)	Spring	29	100.0%	11.1%	3.8%		100.0%	18.8%	0.0%				75.0%					66.7%								
		Fall																								
	Pollock	40	100.0%	100.0%	100.0%		100.0%	100.0%	58.8%		100.0%	100.0%	88.0%		100.0%	100.0%	100.0%		95.5%	18.2%	0.0%					
		23	100.0%	70.0%	0.0%			78.9%	40.4%	0.0%		95.7%	21.5%	0.0%												
Red hake	Spring	20	91.7%	0.0%	0.0%			83.3%	0.0%	0.0%		95.0%	10.0%	0.0%		100.0%	20.0%	0.0%								
		20	91.7%	0.0%	0.0%			83.3%	0.0%	0.0%		95.0%	10.0%	0.0%			100.0%	20.0%	0.0%							
	Fall	20	100.0%	11.1%	0.0%	0.0%		6.7%	0.0%	0.0%	100.0%	35.2%	0.0%	0.0%	100.0%	12.5%	0.0%									
		28	100.0%	88.9%	30.0%	25.0%		93.3%	14.8%	4.0%	100.0%	92.6%	37.0%	2.2%	100.0%	87.5%	0.0%									
Redfish (all years)	Spring	20	100.0%		100.0%	0.0%		100.0%	100.0%		100.0%	100.0%	100.0%							100.0%	100.0%					
		14	100.0%		33.3%	0.0%		100.0%	17.6%		100.0%	90.9%	72.7%							100.0%	50.0%					
	Fall	20					100.0%	100.0%		100.0%	100.0%	100.0%								100.0%	100.0%					
		13					100.0%	92.6%		100.0%	31.1%	100.0%	93.9%	29.5%			100.0%	100.0%	30.7%							
Silver hake	Spring	20	94.6%	16.8%	0.0%		96.8%	31.8%	0.0%		98.6%	40.0%	0.1%		97.7%	44.0%	0.0%									
		19	90.5%	12.7%	0.0%		93.2%	27.2%	0.0%		95.8%	32.6%	0.0%		93.1%	40.7%	0.0%									
	Fall	20	98.7%	22.0%	0.0%	0.0%	100.0%	19.0%	0.0%	0.0%	100.0%	26.7%	0.0%	0.0%	100.0%	15.8%	0.0%	0.0%								
		26	100.0%	87.1%	30.6%	0.0%	100.0%	91.0%	31.2%	5.1%	100.0%	91.4%	28.6%	3.3%	100.0%	86.7%	30.7%	3.0%								
White hake	Spring	25		0.0%	0.0%		4.3%	0.0%	0.0%		26.3%	7.2%	0.0%						13.2%	0.0%						
		34		25.0%	0.0%		78.3%	7.9%	0.0%		90.9%	55.8%	10.8%						83.8%	25.0%						
	Fall	25	50.0%	0.0%	0.0%	77.4%	1.5%	0.0%	0.0%	46.3%	23.5%	0.0%	0.0%	100.0%	49.1%	0.0%	0.0%									
		39	100.0%	25.0%	0.0%	100.0%	82.1%	32.1%	0.0%	100.0%	94.0%	27.0%	0.0%	100.0%	100.0%	43.1%	0.0%									
Winter flounder	Spring	25	100.0%	44.5%	4.5%		100.0%	60.3%	10.3%		100.0%	97.8%	57.2%		100.0%	100.0%	79.3%									
		18	92.6%	4.7%	0.0%		94.3%	6.6%	0.0%		97.5%	40.0%	4.7%		90.0%	44.1%	3.0%									
	Fall	25	88.9%	19.2%	0.0%	100.0%	75.6%	8.6%	0.1%	100.0%	100.0%	77.0%	19.0%		100.0%	96.1%	45.2%									
		28	99.2%	48.0%	4.2%	100.0%	91.8%	25.3%	1.2%	100.0%	100.0%	93.2%	50.4%		100.0%	98.7%	86.3%									
Witch flounder	Spring	30		100.0%	100.0%		100.0%	100.0%	100.0%		100.0%	100.0%	100.0%						100.0%	100.0%						
		20		100.0%	84.0%		100.0%	91.7%	18.2%		100.0%	74.3%	14.7%						100.0%	33.3%	8.6%					
	Fall	30	100.0%	100.0%	100.0%		100.0%	100.0%	100.0%		84.6%	100.0%	100.0%		100.0%	88.0%	100.0%		100.0%	100.0%	66.7%					
		19	100.0%	100.0%	53.3%	11.3%	100.0%	100.0%	0.0%	0.0%	100.0%	88.9%	7.3%	1.4%	100.0%	100.0%	11.1%	0.0%								
Yellowtail flounder	Spring	25	100.0%	23.6%	0.0%		100.0%	19.6%	0.0%		100.0%	62.0%	5.7%						30.0%	5.3%						
		13	96.1%	0.0%	0.0%		92.5%	0.0%	0.0%		67.2%	0.0%	0.0%						0.0%	0.0%						
	Fall	25	100.0%	95.9%	1.5%	0.0%	100.0%	80.8%	0.7%	0.0%	100.0%	91.1%	11.3%	0.3%	100.0%	25.0%	11.1%	0.0%								
		15	73.7%	0.0%	0.0%	0.0%	93.2%	0.2%	0.0%	0.0%	100.0%	1.3%	0.0%	0.0%	100.0%	0.0%	0.0%									

Figure 1. Cod <35 cm distribution by spring survey type, 2002-2012




Table 3. Cumulative number of cod caught by survey over time by size range, compared to 20 percent of total abundance.

COMNAME	ATLANTIC COD							
REGION	(Multiple Items)							
Row Labels	20Pct total num	Num <= 5 cm	Num <= 10 cm	Num <= 15 cm	Num <= 20 cm	Num <= 25 cm	Num <= 30 cm	Num <= 35 cm
IBS Cod Spawning	713	0	1	46	200	309	610	1,340
WINTER	353	0	1	31	99	128	270	737
2002-2012	353	0	1	31	99	128	270	737
SPRING	360	0	0	15	101	181	340	608
2002-2012	360	0	0	15	101	181	340	608
NMFS trawl	19,013	1,824	4,110	6,547	9,888	14,750	22,563	32,232
WINTER	602	2	21	98	247	419	514	599
1963-1971	314	1	20	32	61	118	159	210
1972-1981	92	0	0	0	1	14	22	26
1992-2001	153	1	1	6	34	94	132	162
2002-2012	44	0	0	60	152	194	200	201
SPRING	9,157	1,692	1,815	2,339	3,983	6,797	10,455	14,481
1963-1971	359	62	71	83	104	132	169	259
1972-1981	2,301	443	524	612	901	1,326	1,837	2,507
1982-1991	1,614	78	90	179	396	737	1,113	1,608
1992-2001	607	109	115	141	232	323	427	563
2002-2012	4,276	999	1,015	1,324	2,350	4,280	6,908	9,544
SUMMER	1,486	39	339	440	608	910	1,355	1,849
1963-1971	474	9	18	37	87	118	192	287
1972-1981	847	16	232	282	355	583	905	1,236
1982-1991	23	0	1	1	2	6	14	23
1992-2001	142	14	88	120	159	203	244	302
FALL	7,767	91	1,936	3,670	5,055	6,623	10,240	15,304
1963-1971	660	7	131	190	254	354	523	804
1972-1981	2,140	40	215	356	592	1,045	1,796	2,877
1982-1991	1,111	18	180	299	426	675	1,199	1,847
1992-2001	674	8	57	158	188	272	494	810
2002-2012	3,182	18	1,352	2,667	3,593	4,277	6,228	8,966
MADMF trawl	38,071	136,436	162,095	166,907	172,556	178,159	181,176	183,722
SPRING	32,859	116,635	140,697	143,081	148,325	153,380	155,937	158,110
1972-1981	3,072	4,436	8,954	9,685	11,532	12,971	13,542	14,042
1982-1991	1,382	1,383	1,921	2,440	3,155	4,127	4,824	5,319
1992-2001	3,374	9,900	12,231	12,645	13,756	14,940	15,365	15,831
2002-2012	25,082	100,916	117,592	118,312	119,882	121,342	122,206	122,919
FALL	5,212	19,801	21,398	23,826	24,231	24,779	25,239	25,612
1972-1981	1,874	7,580	7,716	8,444	8,544	8,789	9,006	9,202
1982-1991	1,845	8,230	8,731	8,892	8,981	9,102	9,159	9,201
1992-2001	563	1,124	1,535	2,668	2,709	2,746	2,766	2,787
2002-2012	929	2,867	3,416	3,822	3,996	4,142	4,308	4,423
Grand Total	57,796	138,260	166,206	173,500	182,644	193,218	204,348	217,294

Table 4. Cumulative weight of cod caught by survey over time by size range, compared to 20 percent of total weight.

COMNAME REGION	ATLANTIC COD (Multiple Items)	20 pct total weight	Wgt >= 50 cm	Wgt >= 55 cm	Wgt >= 60 cm	Wgt >= 65 cm	Wgt >= 70 cm	Wgt >= 75 cm	Wgt >= 80 cm	Wgt >= 85 cm	Wgt >= 90 cm
Row Labels											
IBS Cod Spawning		747	2,408	2,110	1,855	1,624	1,347	1,064	798	593	430
WINTER		219	315	184	117	94	50	28	7	7	0
2002-2012		219	315	184	117	94	50	28	7	7	0
SPRING		528	2,093	1,926	1,738	1,530	1,296	1,036	791	587	430
2002-2012		528	2,093	1,926	1,738	1,530	1,296	1,036	791	587	430
NMFS trawl		30,250	126,234	116,874	105,602	91,915	78,010	64,149	52,264	40,675	31,445
WINTER		1,654	7,744	7,421	6,875	6,273	5,663	5,002	4,400	3,594	2,866
1963-1971		1,071	5,112	4,959	4,720	4,403	4,013	3,596	3,173	2,661	2,128
1972-1981		306	1,452	1,397	1,246	1,127	1,070	1,010	923	777	632
1992-2001		269	1,159	1,046	891	724	570	395	305	156	105
2002-2012		8	21	18	18	18	9	2	0	0	0
SPRING		14,558	59,891	55,579	50,284	43,393	36,609	29,872	24,347	18,652	14,302
1963-1971		1,141	5,430	5,229	4,938	4,517	4,148	3,620	3,126	2,501	1,990
1972-1981		4,480	18,878	17,665	16,273	14,448	12,238	10,391	8,984	7,183	5,748
1982-1991		3,639	16,391	15,546	14,307	12,278	10,593	8,661	6,889	5,323	4,055
1992-2001		1,387	6,317	5,887	5,359	4,720	4,063	3,341	2,706	1,977	1,462
2002-2012		3,911	12,875	11,253	9,408	7,430	5,567	3,860	2,642	1,668	1,047
SUMMER		2,879	12,728	11,567	10,206	8,948	7,525	6,234	4,992	3,984	3,132
1963-1971		1,207	5,566	5,241	4,789	4,186	3,500	2,851	2,317	1,769	1,329
1972-1981		1,455	6,301	5,544	4,735	4,162	3,498	2,915	2,279	1,897	1,557
1982-1991		42	172	147	132	104	83	72	68	51	26
1992-2001		174	689	635	550	496	444	395	328	267	220
FALL		11,158	45,872	42,307	38,236	33,302	28,213	23,040	18,526	14,445	11,145
1963-1971		1,684	7,793	7,458	6,993	6,330	5,665	4,982	4,275	3,540	2,821
1972-1981		4,366	19,429	18,092	16,496	14,560	12,593	10,480	8,678	7,073	5,590
1982-1991		1,679	6,914	6,397	5,710	4,879	3,990	3,277	2,553	1,888	1,493
1992-2001		1,063	4,411	3,899	3,322	2,717	2,131	1,512	1,019	702	431
2002-2012		2,365	7,325	6,461	5,716	4,816	3,834	2,789	2,001	1,242	810
MADMF trawl		2,206	5,354	4,219	3,313	2,459	1,767	1,129	736	546	409
SPRING		2,038	5,097	4,015	3,140	2,330	1,681	1,090	715	533	404
1972-1981		407	836	627	445	297	208	149	110	87	71
1982-1991		414	742	533	369	264	180	148	122	101	60
1992-2001		320	633	475	347	225	155	105	60	35	20
2002-2012		896	2,886	2,381	1,980	1,544	1,138	688	423	310	252
FALL		168	257	204	173	130	86	39	22	13	5
1972-1981		61	53	44	37	27	25	16	13	13	5
1982-1991		16	11	9	7	4	3	0	0	0	0
1992-2001		13	12	4	2	0	0	0	0	0	0
2002-2012		78	182	147	126	98	57	23	9	0	0
Grand Total		33,202	133,997	123,202	110,770	95,998	81,124	66,342	53,798	41,814	32,284

Table 5. Cumulative biomass above 5 cm size ranges by species, survey, and decade, compared to 20% of total weight per tow (kg) and the size at estimated 80% maturity for females.

Approximate 20% of biomass (upper), L80 for maturity (lower)	Species	Row Labels	Sum of 20 Pct total wgt																		
			WGT >= 25cm	Wgt >= 30cm	Wgt >= 35cm	Wgt >= 40 cm	Wgt >= 45cm	Wgt >= 50cm	Wgt >= 55 cm	Wgt >= 60cm	Wgt >= 65cm	Wgt >= 70cm	Wgt >= 75cm	Wgt >= 80cm	Wgt >= 85cm	Wgt >= 90cm	Wgt >= 95cm	Wgt >= 100cm			
75 cm	ATLANTIC COD	All	30,250	150,605	149,271	146,284	141,220	134,113	126,234	116,874	105,602	91,915	78,010	64,149	52,264	40,675	31,445	23,602	17,149		
L80 = 50 cm	ATLANTIC COD	WINTER	1,654	8,247	8,226	8,202	8,141	7,983	7,744	7,421	6,875	6,273	5,663	5,002	4,400	3,594	2,866	1,978	1,353		
	ATLANTIC COD	1963-1971	1,071	5,348	5,339	5,325	5,291	5,222	5,112	4,959	4,720	4,403	4,013	3,596	3,173	2,661	2,128	1,461	1,016		
	ATLANTIC COD	1972-1981	306	1,530	1,528	1,527	1,517	1,488	1,452	1,397	1,246	1,127	1,070	1,010	923	777	632	460	312		
	ATLANTIC COD	1992-2001	269	1,339	1,330	1,321	1,305	1,247	1,159	1,046	891	724	570	395	305	156	105	57	25		
	ATLANTIC COD	2002-2012	8	30	29	29	28	26	21	18	18	9	2	0	0	0	0	0	0		
	ATLANTIC COD	SPRING	14,558	72,457	71,801	70,561	68,244	64,198	59,891	55,579	50,284	43,393	36,609	29,872	24,347	18,652	14,302	10,866	7,891		
	ATLANTIC COD	1963-1971	1,141	5,701	5,696	5,672	5,614	5,551	5,430	5,229	4,938	4,517	4,148	3,620	3,126	2,501	1,990	1,516	1,130		
	ATLANTIC COD	1972-1981	4,480	22,342	22,248	22,062	21,645	20,446	18,878	17,665	16,273	14,448	12,238	10,391	8,984	7,183	5,748	4,489	3,320		
	ATLANTIC COD	1982-1991	3,639	18,153	18,082	17,935	17,643	17,118	16,391	15,546	14,307	12,278	10,593	8,661	6,889	5,323	4,055	3,222	2,343		
	ATLANTIC COD	1992-2001	1,387	6,923	6,906	6,864	6,778	6,591	6,317	5,887	4,720	4,063	3,341	2,706	1,977	1,462	1,007	675			
	ATLANTIC COD	2002-2012	3,911	19,338	18,869	18,028	16,564	14,492	12,875	11,253	9,408	7,430	5,567	3,860	2,642	1,668	1,047	632	423		
	ATLANTIC COD	SUMMER	2,879	14,357	14,282	14,124	13,863	13,478	12,728	11,567	10,206	8,948	7,525	6,234	4,992	3,984	3,132	2,334	1,736		
	ATLANTIC COD	1963-1971	1,207	6,032	6,020	5,991	5,927	5,799	5,566	5,241	4,789	4,186	3,500	2,851	2,317	1,769	1,329	974	726		
	ATLANTIC COD	1972-1981	1,455	7,252	7,197	7,088	6,936	6,745	6,301	5,544	4,735	4,162	3,498	2,915	2,279	1,897	1,557	1,169	874		
	ATLANTIC COD	1982-1991	42	209	207	205	203	195	172	147	132	104	83	72	68	51	26	26	26		
	ATLANTIC COD	1992-2001	174	864	858	840	796	739	689	635	550	496	444	395	328	267	220	166	111		
	ATLANTIC COD	FALL	11,158	55,545	54,962	53,397	50,972	48,454	45,872	42,307	38,236	33,302	28,213	23,040	18,526	14,445	11,145	8,424	6,170		
	ATLANTIC COD	1963-1971	1,684	8,407	8,379	8,292	8,177	8,005	7,793	7,458	6,993	6,330	5,665	4,982	4,275	3,540	2,821	2,220	1,622		
	ATLANTIC COD	1972-1981	4,366	21,777	21,653	21,317	20,808	20,197	19,429	18,092	16,496	14,560	12,593	10,480	8,678	7,073	5,590	4,351	3,324		
	ATLANTIC COD	1982-1991	1,679	8,367	8,280	8,078	7,697	7,259	6,914	6,397	5,710	4,879	3,990	3,277	2,553	1,888	1,493	1,046	724		
	ATLANTIC COD	1992-2001	1,063	5,306	5,269	5,173	4,995	4,742	4,411	3,899	3,322	2,717	2,131	1,512	1,019	702	431	293	156		
	ATLANTIC COD	2002-2012	2,365	11,688	11,380	10,536	9,295	8,252	7,325	6,461	5,716	4,816	3,834	2,789	2,001	1,242	810	514	344		
40 cm	AMERICAN PLAICE	WINTER	62	310	300	261	202	130	76	47	22	0	0	0	0	0	0	0	0	0	
L80 = 30 cm	AMERICAN PLAICE	1972-1981	17	85	83	75	63	41	32	27	16	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	1992-2001	44	219	212	182	136	88	44	21	6	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	2002-2012	1	6	5	5	4	1	0	0	0	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	SPRING	2,492	11,176	9,366	6,995	4,939	3,250	1,793	763	289	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	1963-1971	233	1,113	972	756	543	359	194	109	68	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	1972-1981	1,076	4,968	4,453	3,662	2,815	1,951	1,089	482	167	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	1982-1991	453	2,007	1,647	1,216	861	601	366	137	45	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	1992-2001	338	1,498	1,173	757	457	234	105	33	8	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	2002-2012	392	1,589	1,122	603	264	106	38	2	0	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	SUMMER	924	4,013	3,153	2,062	1,264	793	424	171	62	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	1963-1971	81	385	331	244	172	104	65	36	20	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	1972-1981	434	1,875	1,556	1,196	835	544	296	125	38	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	1982-1991	81	350	216	73	20	11	6	0	0	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	1992-2001	328	1,402	1,049	549	237	134	57	11	4	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	FALL	2,690	12,037	10,086	7,423	5,086	3,152	1,750	768	244	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	1963-1971	171	812	706	540	368	224	138	79	39	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	1972-1981	1,248	5,780	5,148	4,197	3,186	2,113	1,221	535	169	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	1982-1991	412	1,777	1,418	982	673	422	234	103	28	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	1992-2001	504	2,217	1,785	1,119	578	265	109	33	8	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	2002-2012	355	1,452	1,030	586	281	128	48	18	0	0	0	0	0	0	0	0	0	0	
	AMERICAN PLAICE	All	6,168	27,535	22,904	16,741	11,491	7,327	4,042	1,750	617	0	0	0	0	0	0	0	0	0	
ATLANTIC HERRING	ATLANTIC HERRING	WINTER	304	765	85	4	2	0	0	0	0	0	0	0	0	0	0	0	0	0	
	ATLANTIC HERRING	1963-1971	8	23	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	ATLANTIC HERRING	1972-1981	9	22	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	ATLANTIC HERRING	1992-2001	260	670	77	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	ATLANTIC HERRING	2002-2012	27	49	2	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	ATLANTIC HERRING	SPRING	2,253	4,363	255	4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	ATLANTIC HERRING	1963-1971	10	23	9	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	ATLANTIC HERRING	1972-1981	239	649	83	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	ATLANTIC HERRING	1982-1991	321	1,063	104	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	ATLANTIC HERRING	1992-2001	778	1,738	46	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	ATLANTIC HERRING	2002-2012	906	890	13	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	ATLANTIC HERRING	SUMMER	1,782	5,508	927	69	2	0	0	0	0	0	0	0	0	0	0	0	0	0	
	ATLANTIC HERRING	1963-1971	229	1,088	615	68	1	0	0	0	0	0	0	0	0	0	0	0	0	0	
	ATLANTIC HERRING	1972-1981	64	220	37	0	0	0	0	0</td											

Approximate 20% of biomass (upper), L80 for maturity (lower)	Species	Row Labels	Sum of 20 Act Total Wgt																	
			Wgt > 25cm	Wgt > 30cm	Wgt > 35cm	Wgt > 40 cm	Wgt > 45cm	Wgt > 50cm	Wgt > 55cm	Wgt > 60cm	Wgt > 65cm	Wgt > 70cm	Wgt > 75cm	Wgt > 80cm	Wgt > 85cm	Wgt > 90cm	Wgt > 95cm	Wgt > 100cm		
50 cm	HADDOCK	All	51,238	243,899	226,195	201,572	172,426	140,490	103,964	68,131	41,692	23,073	11,224	4,337	1,219	0	0	0	0	
L80 = 40 cm	HADDOCK	WINTER	3,338	15,592	14,832	12,926	10,452	8,468	6,770	5,048	3,350	1,972	898	340	66	0	0	0	0	
	HADDOCK	1963-1971	2,933	14,261	13,566	11,708	9,309	7,389	5,820	4,306	2,744	1,578	682	265	49	0	0	0	0	
	HADDOCK	1972-1981	141	707	707	706	703	686	620	471	394	292	168	58	0	0	0	0	0	
	HADDOCK	1992-2001	228	491	432	400	333	291	230	183	136	56	27	11	11	0	0	0	0	
	HADDOCK	2002-2012	35	133	127	112	108	102	99	88	76	46	21	6	6	0	0	0	0	
	HADDOCK	SPRING	16,040	75,439	69,873	59,941	59,644	50,826	38,933	25,459	16,166	9,113	4,571	1,736	510	0	0	0	0	
	HADDOCK	1963-1971	1,416	7,060	7,043	7,001	6,831	6,388	5,732	4,366	2,789	1,492	574	170	27	0	0	0	0	
	HADDOCK	1972-1981	4,819	23,073	22,141	21,302	18,842	15,899	12,933	9,283	6,487	3,895	2,040	796	258	0	0	0	0	
	HADDOCK	1982-1991	1,803	8,905	8,755	8,478	7,793	6,900	5,682	4,175	2,996	1,951	1,150	551	189	0	0	0	0	
	HADDOCK	1992-2001	1,535	7,494	7,330	7,103	6,404	5,589	4,553	3,179	2,049	995	479	121	25	0	0	0	0	
	HADDOCK	2002-2012	6,467	28,907	24,604	22,058	19,774	16,050	10,034	4,455	1,844	780	326	99	11	0	0	0	0	
	HADDOCK	SUMMER	6,262	30,338	27,797	24,468	20,319	14,428	10,562	7,379	4,708	2,538	1,262	478	124	0	0	0	0	
	HADDOCK	1963-1971	4,349	20,828	18,542	15,937	12,591	9,914	7,824	5,390	3,277	1,657	770	209	39	0	0	0	0	
	HADDOCK	1972-1981	1,877	9,338	9,085	8,364	7,570	4,367	2,601	1,872	1,356	844	475	255	85	0	0	0	0	
	HADDOCK	1982-1991	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	HADDOCK	1992-2001	36	172	171	166	158	147	137	117	76	37	18	14	0	0	0	0	0	
	HADDOCK	FALL	25,598	122,530	113,693	98,237	82,011	66,768	47,700	30,245	17,469	9,451	4,493	1,783	519	0	0	0	0	
	HADDOCK	1963-1971	3,186	15,626	15,119	14,014	12,557	10,651	8,649	6,386	4,247	2,411	1,158	401	70	0	0	0	0	
	HADDOCK	1972-1981	6,409	31,571	31,068	27,606	23,347	19,954	15,446	11,065	7,138	4,220	2,278	1,086	343	0	0	0	0	
	HADDOCK	1982-1991	1,664	8,112	7,873	6,994	6,116	5,337	4,397	3,164	1,966	1,114	562	214	88	0	0	0	0	
	HADDOCK	1992-2001	2,978	14,762	14,573	13,737	12,317	10,554	7,506	4,677	2,542	1,235	401	68	18	0	0	0	0	
	HADDOCK	2002-2012	11,361	52,459	45,060	35,885	27,674	20,272	11,703	4,953	1,576	470	94	12	0	0	0	0	0	
100 cm	BARNDOOR SKATE	WINTER	659	3,294	3,292	3,275	3,254	3,211	3,153	3,074	2,991	2,848	2,684	2,535	2,331	2,175	1,995	1,777	1,601	
L80 = 115 cm	BARNDOOR SKATE	1963-1971	207	1,033	1,032	1,032	1,026	1,018	1,001	981	946	895	853	781	720	648	586	559		
	BARNDOOR SKATE	1972-1981	6	29	29	29	29	29	29	29	29	29	29	29	29	29	29	29	29	
	BARNDOOR SKATE	1992-2001	150	750	749	743	736	722	706	682	657	602	566	523	485	458	432	379	331	
	BARNDOOR SKATE	2002-2012	297	1,483	1,483	1,472	1,457	1,435	1,400	1,362	1,324	1,271	1,196	1,131	1,037	969	886	783	682	
	BARNDOOR SKATE	SPRING	495	2,471	2,469	2,463	2,452	2,433	2,401	2,341	2,272	2,171	2,040	1,873	1,765	1,645	1,520	1,420	1,330	
	BARNDOOR SKATE	1963-1971	46	228	228	228	227	226	219	211	193	180	163	163	156	153	143	143	143	
	BARNDOOR SKATE	1972-1981	9	47	47	47	47	47	47	45	43	36	34	34	34	34	21	21	21	
	BARNDOOR SKATE	1982-1991	0	2	2	2	1	1	1	0	0	0	0	0	0	0	0	0	0	
	BARNDOOR SKATE	1992-2001	52	258	258	257	256	254	253	247	240	228	221	211	196	196	192	176	164	
	BARNDOOR SKATE	2002-2012	387	1,936	1,934	1,929	1,920	1,904	1,874	1,831	1,778	1,714	1,606	1,465	1,373	1,260	1,154	1,080	1,001	
	BARNDOOR SKATE	SUMMER	89	443	443	443	443	441	439	433	416	392	361	330	301	265	257	218	183	
	BARNDOOR SKATE	1963-1971	89	443	443	443	443	441	439	433	416	392	361	330	301	265	257	218	183	
	BARNDOOR SKATE	1972-1981	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	BARNDOOR SKATE	1982-1991	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	BARNDOOR SKATE	1992-2001	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	BARNDOOR SKATE	FALL	688	3,438	3,435	3,431	3,421	3,401	3,351	3,279	3,175	3,047	2,919	2,734	2,556	2,410	2,242	2,059	1,893	
	BARNDOOR SKATE	1963-1971	151	756	756	756	756	753	744	731	707	658	614	554	498	446	409	378	334	
	BARNDOOR SKATE	1972-1981	7	34	34	34	34	34	33	30	25	23	23	19	16	16	16	16	16	
	BARNDOOR SKATE	1982-1991	5	26	26	26	26	26	24	23	21	18	14	14	14	10	5	0	0	
	BARNDOOR SKATE	1992-2001	82	410	410	410	409	407	402	397	389	381	374	350	328	303	288	269	263	
	BARNDOOR SKATE	2002-2012	442	2,210	2,208	2,204	2,196	2,181	2,146	2,097	2,031	1,964	1,890	1,797	1,700	1,632	1,518	1,390	1,279	
	BARNDOOR SKATE	All	1,930	9,646	9,639	9,612	9,570	9,486	9,344	9,127	8,854	8,457	8,005	7,472	6,953	6,496	6,014	5,474	5,008	
75 cm	GOOSEFISH	WINTER	1,048	5,221	5,175	5,070	4,902	4,569	4,093	3,474	2,916	2,491	2,014	1,615	1,288	1,019	759	0	0	
L80 = 45 cm	GOOSEFISH	1963-1971	414	2,067	2,064	2,059	2,054	2,036	2,005	1,959	1,882	1,722	1,481	1,256	1,030	848	628	0	0	
	GOOSEFISH	1972-1981	36	179	178	178	177	175	173	171	154	142	123	112	91	75	75	0	0	
	GOOSEFISH	1992-2001	329	1,629	1,599	1,543	1,444	1,245	944	621	440	331	244	151	113	67	56	0	0	
	GOOSEFISH	2002-2012	270	1,346	1,334	1,289	1,226	1,112	971	723	441	296	167	96	54	29	0	0	0	
	GOOSEFISH	SPRING	1,828	9,086	9,024	8,920	8,749	8,487	8,074	7,556	6,979	6,317	5,548	4,716	3,957	3,177	2,449	0	0	
	GOOSEFISH	1963-1971	113	563	562	560	556	557	551	536	511	488	463	389	329	266	159	129	0	
	GOOSEFISH	1972-1981	1,017	5,073	5,050	5,017	4,957	4,863	4,692	4,449	4,187	3,907	3,540	3,083	2,638	2,132	1,615	0	0	
	GOOSEFISH	1982-1991	308	1,537	1,528	1,517	1,500	1,471	1,429	1,364	1,272	1,156	1,011	861	759	694	589	0	0	
	GOOSEFISH	1992-2001	171	833	815	780	732	669	585	506	445	345	286	224	137	96	63	0	0	
	GOOSEFISH	2002-2012	218	1,080	1,068	1,045	1,003	932	831	726	588	446	322	218	157	95	53	0	0	
	GOOSEFISH	SUMMER	646	3,209	3,182	3,140	3,077	3,007	2,923	2,807	2,720	2,563	2,291	2,041	1,778	1,461	1,097	0	0	
	GOOSEFISH	1963-1971	218	1,090	1,090	1,086	1,081	1,069	1,051	1,017	984	921	799	688	576	449	281	0	0	
	GOOSEFISH	1972-1981	334	1,669	1,667	1,664	1,662	1,644	1,631	1,595	1,565	1,492	1,368	1,272	1,121	940	765	0	0	
	GOOSEF																			

Approximate 20% of biomass (upper), L80 for maturity (lower)	Species	Row Labels	Sum of 20 Act Total Wgt																	
			Wgt > 25cm	Wgt > 30cm	Wgt > 35cm	Wgt > 40 cm	Wgt > 45cm	Wgt > 50cm	Wgt > 55cm	Wgt > 60cm	Wgt > 65cm	Wgt > 70cm	Wgt > 75cm	Wgt > 80cm	Wgt > 85cm	Wgt > 90cm	Wgt > 95cm	Wgt > 100cm		
50 cm	LITTLE SKATE	WINTER	4,589	22,768	22,311	21,183	19,260	13,916	2,149	124	34	0	0	0	0	0	0	0	0	
L80 = 55 cm	LITTLE SKATE	1963-1971	457	2,285	2,281	2,257	2,170	1,624	277	32	6	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	1972-1981	144	707	688	637	574	482	221	83	25	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	1992-2001	2,721	13,488	13,186	12,366	11,071	7,779	1,152	8	3	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	2002-2012	1,266	6,288	6,156	5,923	5,444	4,031	498	1	0	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	SPRING	4,842	23,884	23,220	22,036	20,462	16,028	3,493	178	7	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	1963-1971	297	1,476	1,459	1,424	1,360	1,104	239	18	0	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	1972-1981	1,399	6,915	6,758	6,428	5,958	4,685	1,034	74	3	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	1982-1991	1,088	5,359	5,205	4,978	4,665	3,583	795	36	4	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	1992-2001	872	4,277	4,112	3,858	3,554	2,752	604	22	0	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	2002-2012	1,187	5,857	5,686	5,349	4,925	3,905	820	28	0	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	SUMMER	506	2,519	2,505	2,478	2,405	2,005	487	53	5	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	1963-1971	191	951	949	942	918	720	132	30	5	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	1972-1981	271	1,348	1,338	1,320	1,279	1,101	231	4	0	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	1982-1991	0	2	2	2	2	2	0	0	0	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	1992-2001	44	218	217	214	206	182	123	19	0	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	FALL	4,375	21,686	21,347	20,638	19,327	15,447	3,816	213	27	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	1963-1971	342	1,708	1,696	1,666	1,603	1,298	285	41	3	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	1972-1981	1,383	6,853	6,764	6,598	6,256	5,192	1,308	80	16	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	1982-1991	859	4,242	4,137	3,927	3,547	2,701	727	27	0	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	1992-2001	940	4,668	4,604	4,477	4,255	3,403	829	39	7	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	2002-2012	851	4,215	4,145	3,970	3,666	2,853	666	27	0	0	0	0	0	0	0	0	0	0
	LITTLE SKATE	All	14,312	70,856	69,383	66,335	61,454	47,397	9,944	568	73	0	0	0	0	0	0	0	0	0
60 cm	OCEAN POUT	WINTER	1,476	7,370	7,359	7,310	7,176	6,915	6,414	5,599	4,314	2,888	1,919	1,135	584	213	81	0	0	
NA	OCEAN POUT	1963-1971	540	2,700	2,699	2,696	2,689	2,672	2,615	2,459	2,124	1,622	1,219	813	454	177	63	0	0	
	OCEAN POUT	1972-1981	41	203	203	202	200	199	191	168	154	125	83	46	24	8	4	0	0	
	OCEAN POUT	1992-2001	848	4,235	4,225	4,181	4,056	3,823	3,416	2,805	1,909	1,076	575	257	99	29	14	0	0	
	OCEAN POUT	2002-2012	46	232	232	232	231	221	192	166	126	41	20	6	0	0	0	0	0	
	OCEAN POUT	SPRING	2,483	12,390	12,343	12,201	11,861	11,029	9,865	8,242	6,549	4,631	3,047	1,720	904	381	137	0	0	
	OCEAN POUT	1963-1971	146	728	728	725	718	684	607	549	467	370	283	159	94	41	24	0	0	
	OCEAN POUT	1972-1981	710	3,541	3,527	3,484	3,363	2,974	2,517	2,010	1,575	1,128	743	455	281	125	43	0	0	
	OCEAN POUT	1982-1991	1,111	5,546	5,529	5,473	5,343	5,078	4,685	3,986	3,196	2,271	1,468	829	410	175	67	0	0	
	OCEAN POUT	1992-2001	353	1,764	1,759	1,742	1,706	1,621	1,471	1,209	914	598	392	201	89	33	3	0	0	
	OCEAN POUT	2002-2012	163	810	801	776	732	671	585	489	397	264	162	76	31	6	0	0	0	
	OCEAN POUT	SUMMER	277	1,384	1,375	1,345	1,277	1,170	1,042	918	787	629	453	273	146	55	26	0	0	
	OCEAN POUT	1963-1971	95	473	472	471	466	459	452	439	407	340	236	128	62	28	12	0	0	
	OCEAN POUT	1972-1981	127	631	625	608	578	531	456	396	329	269	203	143	84	28	13	0	0	
	OCEAN POUT	1982-1991	15	73	72	70	62	46	32	22	13	10	8	2	0	0	0	0	0	
	OCEAN POUT	1992-2001	42	207	205	197	171	134	101	62	38	10	6	0	0	0	0	0	0	
	OCEAN POUT	FALL	446	2,216	2,188	2,088	1,908	1,663	1,358	1,027	729	481	293	183	114	59	28	0	0	
	OCEAN POUT	1963-1971	54	271	269	264	251	231	205	166	137	104	60	38	25	11	11	0	0	
	OCEAN POUT	1972-1981	151	752	744	725	686	620	526	404	291	185	137	97	63	40	13	0	0	
	OCEAN POUT	1982-1991	85	422	416	395	364	315	243	182	119	77	49	23	13	4	4	0	0	
	OCEAN POUT	1992-2001	111	552	546	523	465	395	312	233	158	102	45	25	13	4	0	0	0	
	OCEAN POUT	2002-2012	45	219	212	182	142	102	72	42	25	14	1	0	0	0	0	0	0	
	OCEAN POUT	All	4,682	23,360	23,265	22,943	22,221	20,777	18,679	15,786	12,378	8,629	5,712	3,311	1,748	707	273	0	0	
75 cm	POLLOCK	WINTER	621	3,094	3,071	3,039	2,934	2,838	2,712	2,576	2,384	2,143	1,800	1,466	1,051	607	311	139	0	
L80 = 50 cm	POLLOCK	1963-1971	505	2,518	2,495	2,463	2,359	2,266	2,142	2,013	1,845	1,630	1,351	1,094	761	416	195	89	0	
	POLLOCK	1972-1981	106	529	529	528	528	525	523	517	498	473	413	340	273	174	105	40	0	
	POLLOCK	1992-2001	10	48	48	48	47	47	47	45	41	39	36	32	17	17	10	10	0	
	POLLOCK	2002-2012	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	POLLOCK	SPRING	5,183	25,770	25,582	25,096	24,484	23,329	22,026	20,190	17,838	15,673	13,483	11,170	8,798	6,597	4,321	2,219	0	
	POLLOCK	1963-1971	459	2,286	2,280	2,270	2,257	2,233	2,194	2,158	2,077	1,996	1,964	1,859	1,608	1,166	632	244	0	
	POLLOCK	1972-1981	1,753	8,743	8,651	8,337	8,009	7,547	7,201	6,720	6,088	5,590	5,054	4,547	3,889	3,065	1,997	1,040	0	
	POLLOCK	1982-1991	1,630	8,125	8,093	8,038	7,951	7,600	6,981	6,114	5,196	4,547	3,650	2,950	2,385	1,964	1,481	851	0	
	POLLOCK	1992-2001	513	2,533	2,500	2,448	2,305	2,036	1,818	1,589	1,351	1,079	864	643	377	170	100	44	0	
	POLLOCK	2002-2012	828	4,084	4,058	4,003	3,961	3,914	3,833	3,609	3,126	2,551	1,951	1,171	540	232	110	40	0	
	POLLOCK	SUMMER	812	3,975	3,913	3,881	3,805	3,705	3,616	3,459	3,285	3,089	2,738	2,273	1,797	1,298	820	458	0	
	POLLOCK	1963-1971	349	1,747	1,746	1,735	1,694	1,614	1,538	1,427	1,343	1,244	1,093	847	575	304	132	48	0	
	POLLOCK	1972-1981	429	2,076	2,025	2,012	1,982	1,964	1,950	1,909	1,827	1,745	1,578	1,395	1,204	976	677	399	0	
	POLLOCK	1982-1991	1	5	4	4	3	2	2	0	0	0	0	0	0	0	0	0	0	
	POLLOCK	1992-2001	33	147	138	131	126	125	125	121	115	100	67	32	19	19	11	11	0	
	POLLOCK	FALL	4,206	20,730																

Approximate 20% of biomass (upper), L80 for maturity (lower)	Species	Row Labels	Sum of 20 pct Total Wgt																
			WGT >= 25cm	Wgt >= 30cm	Wgt >= 35cm	Wgt >= 40 cm	Wgt >= 45cm	Wgt >= 50cm	Wgt >= 55cm	Wgt >= 60cm	Wgt >= 65cm	Wgt >= 70cm	Wgt >= 75cm	Wgt >= 80cm	Wgt >= 85cm	Wgt >= 90cm	Wgt >= 95cm	Wgt >= 100cm	
35 cm	RED HAKE	WINTER	818	3,968	2,731	1,249	497	199	83	30	0	0	0	0	0	0	0	0	0
L80 = 35 cm	RED HAKE	1963-1971	317	1,570	1,470	892	395	178	77	29	0	0	0	0	0	0	0	0	0
	RED HAKE	1972-1981	2	11	11	7	4	2	2	2	0	0	0	0	0	0	0	0	0
	RED HAKE	1992-2001	419	2,046	1,185	336	96	19	4	0	0	0	0	0	0	0	0	0	0
	RED HAKE	2002-2012	80	341	67	15	2	0	0	0	0	0	0	0	0	0	0	0	0
	RED HAKE	SPRING	2,156	10,414	8,692	5,260	2,749	1,180	438	128	0	0	0	0	0	0	0	0	0
	RED HAKE	1963-1971	80	393	367	257	139	69	32	10	0	0	0	0	0	0	0	0	0
	RED HAKE	1972-1981	718	3,501	3,185	2,149	1,224	562	237	78	0	0	0	0	0	0	0	0	0
	RED HAKE	1982-1991	427	2,066	1,871	1,298	737	337	117	33	0	0	0	0	0	0	0	0	0
	RED HAKE	1992-2001	427	2,064	1,662	912	435	156	40	4	0	0	0	0	0	0	0	0	0
	RED HAKE	2002-2012	504	2,390	1,607	644	214	56	13	3	0	0	0	0	0	0	0	0	0
	RED HAKE	SUMMER	825	4,045	3,797	2,714	1,508	667	249	88	0	0	0	0	0	0	0	0	0
	RED HAKE	1963-1971	191	928	858	562	282	135	56	21	0	0	0	0	0	0	0	0	0
	RED HAKE	1972-1981	383	1,887	1,817	1,357	790	355	143	54	0	0	0	0	0	0	0	0	0
	RED HAKE	1982-1991	30	147	143	109	59	33	9	1	0	0	0	0	0	0	0	0	0
	RED HAKE	1992-2001	221	1,083	980	686	376	144	41	12	0	0	0	0	0	0	0	0	0
	RED HAKE	FALL	3,613	17,177	14,333	9,416	4,954	2,143	744	223	0	0	0	0	0	0	0	0	0
	RED HAKE	1963-1971	257	1,246	1,113	786	403	200	75	15	0	0	0	0	0	0	0	0	0
	RED HAKE	1972-1981	1,087	5,270	4,767	3,458	1,895	844	322	107	0	0	0	0	0	0	0	0	0
	RED HAKE	1982-1991	762	3,681	3,152	2,259	1,314	633	225	68	0	0	0	0	0	0	0	0	0
	RED HAKE	1992-2001	838	3,878	3,129	1,919	979	376	99	24	0	0	0	0	0	0	0	0	0
	RED HAKE	2002-2012	670	3,102	2,172	994	363	90	22	10	0	0	0	0	0	0	0	0	0
	RED HAKE	All	7,413	35,603	29,554	18,639	9,708	4,190	1,514	469	0	0	0	0	0	0	0	0	0
30 cm	ACADIAN REDFISH	WINTER	745	3,127	1,895	705	35	3	0	0	0	0	0	0	0	0	0	0	0
L80 = 25 cm	ACADIAN REDFISH	1963-1971	745	3,127	1,895	705	35	3	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	1972-1981	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	1992-2001	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	2002-2012	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	SPRING	9,999	40,176	23,508	8,686	1,887	307	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	1963-1971	1,010	4,384	3,038	1,333	190	13	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	1972-1981	2,415	11,202	8,598	4,513	1,259	269	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	1982-1991	646	3,049	2,471	1,219	292	20	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	1992-2001	2,212	6,703	3,099	687	94	2	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	2002-2012	3,716	14,838	6,303	934	52	4	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	SUMMER	2,449	10,299	6,804	2,913	463	28	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	1963-1971	1,859	8,020	5,280	2,060	274	23	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	1972-1981	401	1,787	1,298	779	170	5	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	1982-1991	12	29	13	5	0	0	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	1992-2001	178	464	212	69	18	0	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	FALL	14,447	57,004	33,362	12,479	2,454	95	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	1963-1971	2,272	9,463	6,746	2,739	349	19	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	1972-1981	2,895	13,232	10,478	5,990	1,477	46	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	1982-1991	865	3,812	2,990	1,545	382	12	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	1992-2001	2,188	6,929	2,720	727	167	8	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	2002-2012	6,227	23,569	10,428	1,478	80	10	0	0	0	0	0	0	0	0	0	0	0
	ACADIAN REDFISH	All	27,641	110,606	65,569	24,782	4,839	433	0	0	0	0	0	0	0	0	0	0	0
	ROSETTE SKATE	WINTER	6	31	30	29	24	0	0	0	0	0	0	0	0	0	0	0	0
L80 = 55 cm	ROSETTE SKATE	1963-1971	0	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0
	ROSETTE SKATE	1972-1981	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	ROSETTE SKATE	1992-2001	2	12	12	12	11	0	0	0	0	0	0	0	0	0	0	0	0
	ROSETTE SKATE	2002-2012	4	18	18	17	13	0	0	0	0	0	0	0	0	0	0	0	0
	ROSETTE SKATE	SPRING	1	3	3	2	0	0	0	0	0	0	0	0	0	0	0	0	0
	ROSETTE SKATE	1963-1971	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	ROSETTE SKATE	1972-1981	0	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0
	ROSETTE SKATE	1982-1991	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	ROSETTE SKATE	1992-2001	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	ROSETTE SKATE	2002-2012	0	2	2	2	0	0	0	0	0	0	0	0	0	0	0	0	0
	ROSETTE SKATE	SUMMER	0	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0
	ROSETTE SKATE	1963-1971	0	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0
	ROSETTE SKATE	1972-1981	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	ROSETTE SKATE	1982-1991	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	ROSETTE SKATE	1992-2001	0	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0
	ROSETTE SKATE	2002-2012	3	16	15	14	7	0	0	0	0	0	0	0	0	0	0	0	0
	ROSETTE SKATE	All	11	54	53	50	37	0	0	0	0	0	0	0	0	0	0	0	0

Approximate 20% of biomass (upper), L80 for maturity (lower)	Species	Row Labels	Sum of 20 Act Total Wgt																	
			Wgt > 25cm	Wgt > 30cm	Wgt > 35cm	Wgt > 40 cm	Wgt > 45cm	Wgt > 50cm	Wgt > 55cm	Wgt > 60cm	Wgt > 65cm	Wgt > 70cm	Wgt > 75cm	Wgt > 80cm	Wgt > 85cm	Wgt > 90cm	Wgt > 95cm	Wgt > 100cm		
30 cm	SILVER HAKE	WINTER	530	1,815	675	312	134	78	44	13	0	0	0	0	0	0	0	0	0	
L80 = 30 cm	SILVER HAKE	1963-1971	208	775	443	241	108	64	40	11	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	1972-1981	4	19	15	9	7	6	3	2	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	1992-2001	280	919	185	51	17	6	1	0	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	2002-2012	39	102	33	11	3	2	0	0	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	SPRING	3,994	12,959	6,550	2,564	1,024	508	284	152	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	1963-1971	70	298	189	102	49	26	8	2	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	1972-1981	1,714	6,911	4,682	1,876	727	381	219	115	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	1982-1991	484	1,678	789	289	118	52	30	18	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	1992-2001	1,045	2,517	486	183	90	33	20	13	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	2002-2012	681	1,555	404	114	40	16	6	4	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	SUMMER	1,639	5,840	3,990	1,837	853	467	277	125	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	1963-1971	571	2,651	1,873	821	354	184	114	50	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	1972-1981	438	1,927	1,579	807	414	242	135	64	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	1982-1991	94	206	108	42	9	7	4	0	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	1992-2001	535	1,056	430	167	75	34	24	11	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	FALL	6,532	23,582	13,035	5,751	2,586	1,322	727	364	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	1963-1971	569	2,436	1,754	911	528	339	198	94	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	1972-1981	1,417	6,111	4,801	2,432	1,091	630	401	222	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	1982-1991	1,525	6,284	3,577	1,470	577	189	55	26	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	1992-2001	1,530	4,656	1,738	554	243	112	46	14	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	2002-2012	1,491	4,093	1,167	384	148	53	27	8	0	0	0	0	0	0	0	0	0	
	SILVER HAKE	All	12,695	44,196	24,250	10,463	4,597	2,376	1,332	654	0	0	0	0	0	0	0	0	0	
55 cm	SMOOTH SKATE	WINTER	33	165	162	154	142	128	109	67	18	0	0	0	0	0	0	0	0	
L80 = 65 cm	SMOOTH SKATE	1963-1971	16	78	76	72	66	60	52	29	7	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	1972-1981	10	52	50	47	43	39	34	24	5	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	1992-2001	7	35	35	34	33	29	23	14	5	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	2002-2012	0	1	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	SPRING	226	1,115	1,095	1,057	995	900	712	382	109	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	1963-1971	23	116	115	113	108	103	91	54	18	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	1972-1981	77	382	376	365	344	309	250	141	46	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	1982-1991	35	172	169	165	159	149	127	74	27	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	1992-2001	25	124	122	116	112	102	75	36	4	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	2002-2012	66	322	313	298	272	236	168	76	15	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	SUMMER	26	129	127	124	118	107	90	55	17	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	1963-1971	12	58	58	57	56	51	42	26	10	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	1972-1981	5	27	27	26	25	21	18	10	3	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	1982-1991	2	12	11	11	9	9	8	4	0	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	1992-2001	7	32	31	30	28	26	22	15	3	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	FALL	247	1,219	1,199	1,166	1,118	1,041	892	511	152	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	1963-1971	39	191	188	182	173	162	141	82	22	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	1972-1981	58	291	289	285	278	261	223	124	43	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	1982-1991	39	195	192	189	182	173	154	97	34	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	1992-2001	55	272	266	257	246	223	187	104	28	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	2002-2012	56	271	264	253	240	222	188	105	25	0	0	0	0	0	0	0	0	
	SMOOTH SKATE	All	532	2,628	2,583	2,502	2,373	2,176	1,804	1,015	296	0	0	0	0	0	0	0	0	
85 cm	THORNY SKATE	WINTER	592	2,945	2,927	2,893	2,852	2,795	2,723	2,614	2,482	2,320	2,130	1,920	1,640	1,205	854	468	189	
L80 = 95 cm	THORNY SKATE	1963-1971	486	2,422	2,410	2,389	2,368	2,334	2,291	2,218	2,123	2,005	1,864	1,685	1,467	1,130	829	450	189	
	THORNY SKATE	1972-1981	83	413	409	404	395	382	362	339	313	280	243	215	158	69	25	18	0	
	THORNY SKATE	1992-2001	22	109	107	98	87	76	69	56	46	35	23	20	16	6	0	0	0	
	THORNY SKATE	2002-2012	0	2	2	2	2	2	2	0	0	0	0	0	0	0	0	0	0	
	THORNY SKATE	SPRING	2,268	11,258	11,162	11,035	10,829	10,557	10,115	9,495	8,737	7,931	7,090	6,159	5,186	4,047	2,771	1,691	869	
	THORNY SKATE	1963-1971	474	2,354	2,338	2,324	2,295	2,250	2,166	2,094	1,979	1,871	1,710	1,556	1,371	1,094	779	494	290	
	THORNY SKATE	1972-1981	1,059	5,262	5,223	5,162	5,068	4,944	4,757	4,448	4,088	3,683	3,288	2,801	2,353	1,914	1,280	833	450	
	THORNY SKATE	1982-1991	495	2,459	2,435	2,406	2,355	2,297	2,207	2,057	1,881	1,660	1,460	1,256	1,013	721	508	279	96	
	THORNY SKATE	1992-2001	134	663	654	643	625	599	556	510	446	397	353	309	254	168	103	41	22	
	THORNY SKATE	2002-2012	105	520	512	501	486	468	429	385	344	320	279	237	195	149	102	43	10	
	THORNY SKATE	SUMMER	952	4,741	4,719	4,687	4,642	4,576	4,483	4,330	4,095	3,821	3,498	3,089	2,636	2,053	1,528	847	321	
	THORNY SKATE	1963-1971	527	2,627	2,617	2,607	2,587	2,554	2,504	2,437	2,329	2,199	2,050	1,862	1,627	1,324	1,086	660	275	
	THORNY SKATE	1972-1981	315	1,570	1,566	1,553	1,539	1,515	1,493	1,440	1,354	1,255	1,119	934	772	562	339	152	45	
	THORNY SKATE	1982-1991	35	174	171	169	168	165	160	157	150	146	134	116	91	64	31	9	0	
	THORNY SKATE	1992-2001	75	369	364	359	349	342	325	296	262	221	195	177	147	104	72	26	0	
	THORNY SKATE	FALL	1,069	18,154	18,050	17,923	17,842	16,831	16,930	16,137	15,760	15,420	15,169	14,884	14,544	6,884	4,528	2,952	1,212	
	THORNY SKATE	1963-1971	1,141	5,679	5,651	5,609	5,559	5,484	5,392	5,245	5,032	4,760	4,461	4,037	3,575	2,969	2,339	1,565	691	
	THORNY SKATE	1972-1981	489	2,427	2,408	2,379	2,329	2,268	2,172	2,049	1,866	1,695	1,482	1,244	1,023	745	535	326	160	
	THORNY SKATE	1982-1991	284	1,408	1,396	1,377	1,351	1,303	1,230	1,139	1,0									

Approximate 20% of biomass (upper), L80 for maturity (lower)	Species	Row Labels	Sum of 20 Act Total Wgt																		
			Wgt > 25cm	Wgt > 30cm	Wgt > 35cm	Wgt > 40 cm	Wgt > 45cm	Wgt > 50cm	Wgt > 55cm	Wgt > 60cm	Wgt > 65cm	Wgt > 70cm	Wgt > 75cm	Wgt > 80cm	Wgt > 85cm	Wgt > 90cm	Wgt > 95cm	Wgt > 100cm			
75 cm	WHITE HAKE	WINTER	302	1,502	1,483	1,427	1,349	1,248	1,134	1,051	955	813	639	515	445	397	352	313	295		
L80 = 45 cm	WHITE HAKE	1963-1971	258	1,286	1,270	1,247	1,194	1,107	1,024	952	878	755	609	491	421	378	339	300	282		
	WHITE HAKE	1972-1981	18	90	90	79	71	69	54	49	40	31	16	13	13	13	13	13	13	13	
	WHITE HAKE	1992-2001	19	93	90	74	61	53	43	38	28	21	14	11	11	6	0	0	0	0	
	WHITE HAKE	2002-2012	7	33	33	27	23	20	14	11	8	5	0	0	0	0	0	0	0	0	
	WHITE HAKE	SPRING	3,694	18,429	18,187	17,524	16,803	15,598	14,114	12,786	11,344	9,412	7,425	5,441	3,983	2,905	2,405	1,950	1,581		
	WHITE HAKE	1963-1971	170	849	839	816	769	690	614	561	506	432	364	321	273	240	212	171	138		
	WHITE HAKE	1972-1981	1,691	8,445	8,358	8,125	7,843	7,410	6,813	6,296	5,769	5,008	4,198	3,157	2,331	1,610	1,320	1,118	961		
	WHITE HAKE	1982-1991	795	3,967	3,900	3,712	3,538	3,270	2,966	2,698	2,346	1,919	1,413	981	695	572	494	422	356		
	WHITE HAKE	1992-2001	450	2,246	2,211	2,115	2,014	1,802	1,523	1,289	1,088	786	523	339	210	148	121	84	44		
	WHITE HAKE	2002-2012	587	2,923	2,879	2,756	2,639	2,425	2,198	1,942	1,636	1,267	927	643	475	334	259	155	82		
	WHITE HAKE	SUMMER	1,171	5,840	5,741	5,426	4,997	4,494	3,956	3,489	3,087	2,507	1,885	1,381	1,013	719	587	504	437		
	WHITE HAKE	1963-1971	355	1,776	1,770	1,745	1,700	1,614	1,515	1,417	1,300	1,088	822	566	426	333	272	236	204		
	WHITE HAKE	1972-1981	414	2,070	2,062	1,998	1,861	1,722	1,561	1,416	1,290	1,089	884	715	537	369	316	268	233		
	WHITE HAKE	1982-1991	135	672	652	562	436	343	247	174	124	73	32	20	9	0	0	0	0		
	WHITE HAKE	1992-2001	266	1,322	1,257	1,121	1,000	815	633	482	374	258	147	80	40	16	0	0	0		
	WHITE HAKE	FALL	5,519	27,377	26,873	26,313	24,673	22,062	19,488	17,049	14,531	11,918	9,129	6,826	5,143	3,764	2,940	2,370	1,933		
	WHITE HAKE	1963-1971	779	3,885	3,826	3,725	3,542	3,217	2,909	2,616	2,284	1,899	1,509	1,136	897	716	651	528	490		
	WHITE HAKE	1972-1981	2,231	11,109	10,951	10,783	10,258	9,366	8,471	7,547	6,702	5,769	4,647	3,640	2,803	2,033	1,654	1,371	1,151		
	WHITE HAKE	1982-1991	1,080	5,307	5,164	5,020	4,548	3,881	3,308	2,822	2,313	1,840	1,354	960	628	402	243	182	142		
	WHITE HAKE	1992-2001	801	3,968	3,891	3,798	3,537	3,120	2,646	2,188	1,705	1,237	788	533	412	329	231	168	73		
	WHITE HAKE	2002-2012	628	3,108	3,042	2,988	2,787	2,478	2,154	1,876	1,527	1,173	830	558	404	284	162	120	76		
	WHITE HAKE	All	10,687	53,149	52,284	50,691	47,823	43,402	38,693	34,375	29,917	24,650	19,078	14,164	10,583	7,784	6,285	5,138	4,247		
30 cm	WINDOWPANE	WINTER	1,033	4,331	3,104	119	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
L80 = 25 cm	WINDOWPANE	1963-1971	28	134	77	13	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	1972-1981	15	66	44	13	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	1992-2001	869	3,573	978	79	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	2002-2012	121	557	205	14	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	SPRING	834	3,681	1,863	426	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	1963-1971	20	91	51	8	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	1972-1981	439	1,948	948	186	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	1982-1991	238	1,074	638	211	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	1992-2001	75	306	124	15	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	2002-2012	62	262	102	6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	SUMMER	101	484	327	76	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	1963-1971	19	94	67	7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	1972-1981	81	387	260	69	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	1982-1991	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	1992-2001	1	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	FALL	1,097	4,636	2,200	420	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	1963-1971	54	230	109	19	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	1972-1981	370	1,668	955	200	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	1982-1991	251	1,055	607	157	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	1992-2001	263	1,077	374	35	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	2002-2012	159	607	155	10	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINDOWPANE	All	3,066	13,132	5,695	1,041	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
45 cm	WINTER FLOUNDER	WINTER	271	1,340	1,287	1,140	910	620	316	126	15	3	0	0	0	0	0	0	0	0	
L80 = 30 cm	WINTER FLOUNDER	1963-1971	157	782	767	718	600	415	192	78	12	3	0	0	0	0	0	0	0	0	
	WINTER FLOUNDER	1972-1981	43	214	209	188	165	132	87	40	3	0	0	0	0	0	0	0	0	0	
	WINTER FLOUNDER	1992-2001	57	278	250	183	115	55	27	9	0	0	0	0	0	0	0	0	0	0	
	WINTER FLOUNDER	2002-2012	14	67	61	50	31	17	10	0	0	0	0	0	0	0	0	0	0	0	
	WINTER FLOUNDER	SPRING	2,113	9,986	8,765	6,791	4,642	2,690	1,090	344	94	11	0	0	0	0	0	0	0	0	
	WINTER FLOUNDER	1963-1971	149	739	722	686	551	382	202	52	14	0	0	0	0	0	0	0	0	0	
	WINTER FLOUNDER	1972-1981	650	3,164	2,906	2,392	1,698	1,003	431	169	53	3	0	0	0	0	0	0	0	0	
	WINTER FLOUNDER	1982-1991	551	2,606	2,312	1,788	1,193	626	220	65	21	7	0	0	0	0	0	0	0	0	
	WINTER FLOUNDER	1992-2001	279	1,323	1,161	834	535	271	96	23	0	0	0	0	0	0	0	0	0	0	
	WINTER FLOUNDER	2002-2012	484	2,154	1,663	1,092	665	408	141	34	5	0	0	0	0	0	0	0	0	0	
	WINTER FLOUNDER	SUMMER	799	3,690	3,069	2,101	1,314	693	349	154	38	3	0	0	0	0	0	0	0	0	
	WINTER FLOUNDER	1963-1971	159	794	776	709	564	305	140	62	18	3	0	0	0	0	0	0	0	0	
	WINTER FLOUNDER	1972-1981	529	2,437	1,978	1,274	709	382	208	92	20	0	0	0	0	0	0	0	0	0	
	WINTER FLOUNDER	1982-1991	6	25	16	8	2	0	0	0	0	0	0	0	0	0	0	0	0	0	
	WINTER FLOUNDER	1992-2001	105	434	300	110	39	6	0	0	0	0	0	0	0	0	0	0	0	0	
	WINTER FLOUNDER	Fall	1,114	14,652	12,372	5,244	5,740	3,254	1,588	588	153	31	0	1	0	0	0	0	0	0	
	WINTER FLOUNDER	1963-1971	234	1,165	1,136	1,064	895	611	348	169	66	23	0	0	0	0	0	0	0	0	
	WINTER FLOUNDER	1972-1981																			

Approximate 20% of biomass (upper), L80 for maturity (lower)	Species	Row Labels	Sum of 20 Act Total Wgt																		
			Wgt > 25cm	Wgt > 30cm	Wgt > 35cm	Wgt > 40 cm	Wgt > 45cm	Wgt > 50cm	Wgt > 55cm	Wgt > 60cm	Wgt > 65cm	Wgt > 70cm	Wgt > 75cm	Wgt > 80cm	Wgt > 85cm	Wgt > 90cm	Wgt > 95cm	Wgt > 100cm			
85 cm	WINTER SKATE	WINTER	4,668	23,318	23,259	22,960	22,031	20,363	18,249	16,314	14,709	13,096	11,643	9,629	7,502	5,206	3,107	1,294	0		
L80 = 95 cm	WINTER SKATE	1963-1971	383	1,913	1,913	1,910	1,891	1,815	1,599	1,377	1,251	1,145	1,070	925	752	543	313	153	0		
	WINTER SKATE	1972-1981	262	1,312	1,311	1,307	1,296	1,273	1,243	1,179	1,122	1,007	903	704	418	244	112	44	0		
	WINTER SKATE	1992-2001	2,655	13,268	13,234	12,982	12,226	10,878	9,282	7,891	6,776	5,746	4,860	3,816	2,991	2,132	1,446	827	0		
	WINTER SKATE	2002-2012	1,368	6,824	6,802	6,760	6,618	6,397	6,126	5,867	5,560	5,198	4,811	4,185	3,341	2,286	1,236	270	0		
	WINTER SKATE	SPRING	9,956	49,756	49,672	49,296	48,195	46,627	44,769	42,691	40,306	37,361	34,054	29,903	24,996	18,536	12,538	7,691	0		
	WINTER SKATE	1963-1971	390	1,949	1,948	1,945	1,928	1,891	1,809	1,685	1,480	1,239	1,005	727	544	390	232	143	0		
	WINTER SKATE	1972-1981	1,357	6,783	6,776	6,753	6,686	6,593	6,480	6,283	6,024	5,661	5,132	4,454	3,557	2,387	1,305	736	0		
	WINTER SKATE	1982-1991	5,405	27,006	26,950	26,715	26,134	25,429	24,699	23,936	23,122	22,029	20,807	19,070	16,886	13,331	9,715	6,253	0		
	WINTER SKATE	1992-2001	1,238	6,187	6,180	6,132	5,921	5,490	4,899	4,333	3,764	3,172	2,612	2,105	1,572	1,004	631	373	0		
	WINTER SKATE	2002-2012	1,567	7,832	7,819	7,751	7,225	6,888	6,454	5,916	5,260	4,497	3,547	2,437	1,424	655	185	0			
	WINTER SKATE	SUMMER	1,968	9,839	9,836	9,821	9,780	9,693	9,524	9,339	9,120	8,869	8,522	7,951	6,903	5,141	3,275	1,814	0		
	WINTER SKATE	1963-1971	318	1,589	1,588	1,586	1,580	1,551	1,459	1,342	1,217	1,102	964	784	589	394	233	114	0		
	WINTER SKATE	1972-1981	1,633	8,163	8,162	8,152	8,124	8,071	7,997	7,933	7,844	7,716	7,514	7,128	6,285	4,735	3,035	1,699	0		
	WINTER SKATE	1982-1991	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	WINTER SKATE	1992-2001	17	86	86	83	75	71	68	64	60	51	44	39	29	12	6	0	0		
	WINTER SKATE	FALL	13,916	69,553	69,461	69,078	68,009	66,538	64,471	61,888	58,517	54,479	49,958	44,159	36,580	26,821	17,513	10,275	0		
	WINTER SKATE	1963-1971	431	2,151	2,146	2,126	2,096	2,020	1,852	1,678	1,453	1,234	1,028	810	589	412	347	233	0		
	WINTER SKATE	1972-1981	2,861	14,300	14,276	14,209	14,065	13,848	13,580	13,290	12,953	12,471	11,813	10,649	8,916	6,472	4,068	2,342	0		
	WINTER SKATE	1982-1991	4,979	24,882	24,842	24,731	24,468	24,113	23,667	23,194	22,527	21,721	20,782	19,427	17,664	14,260	9,876	6,158	0		
	WINTER SKATE	1992-2001	2,415	12,069	12,059	12,010	11,823	11,453	10,773	8,573	7,222	5,984	4,843	3,587	2,396	1,543	910	0	0		
	WINTER SKATE	2002-2012	3,231	16,151	16,138	16,002	15,557	15,105	14,600	13,956	13,010	11,831	10,352	8,429	5,823	3,280	1,678	632	0		
	WINTER SKATE	All	30,508	152,466	152,229	151,154	148,015	143,222	137,014	130,231	122,652	113,805	104,177	91,643	75,982	55,703	36,432	21,074	0		
45 cm	WITCH FLOUNDER	WINTER	217	1,079	1,018	951	788	545	336	181	41	0	0	0	0	0	0	0	0		
L80 = 40 cm	WITCH FLOUNDER	1963-1971	118	586	582	564	526	441	319	178	40	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	1972-1981	2	9	9	9	9	7	3	0	0	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	1992-2001	54	271	269	255	185	71	11	3	2	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	2002-2012	43	213	158	123	69	26	3	0	0	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	SPRING	997	4,916	4,748	4,332	3,715	3,006	2,039	926	186	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	1963-1971	140	697	692	674	636	528	324	147	38	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	1972-1981	508	2,511	2,457	2,328	2,118	1,854	1,320	589	103	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	1982-1991	153	757	735	684	602	482	348	172	42	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	1992-2001	70	334	297	220	123	68	33	15	3	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	2002-2012	126	618	568	426	235	75	15	2	0	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	SUMMER	278	1,356	1,314	1,224	1,092	925	690	366	94	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	1963-1971	129	642	635	616	554	456	324	182	44	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	1972-1981	107	530	522	505	472	423	334	169	42	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	1982-1991	11	48	43	31	20	15	10	4	2	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	1992-2001	31	135	114	72	46	30	22	11	6	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	FALL	980	4,842	4,663	4,294	3,750	3,055	2,176	1,093	267	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	1963-1971	286	1,427	1,413	1,368	1,263	1,041	719	334	90	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	1972-1981	405	2,012	1,969	1,895	1,784	1,579	1,183	604	123	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	1982-1991	106	526	507	466	402	314	225	140	53	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	1992-2001	97	460	390	271	153	74	38	14	1	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	2002-2012	86	418	384	293	148	47	10	0	0	0	0	0	0	0	0	0	0		
	WITCH FLOUNDER	All	2,472	12,193	11,744	10,800	9,345	7,530	5,241	2,566	589	0	0	0	0	0	0	0	0		
40 cm	YELLOWTAIL FLOUNDER	WINTER	1,267	6,287	5,679	3,978	1,812	394	49	0	0	0	0	0	0	0	0	0	0		
L80 = 30 cm	YELLOWTAIL FLOUNDER	1963-1971	213	1,028	958	767	406	116	18	0	0	0	0	0	0	0	0	0	0		
	YELLOWTAIL FLOUNDER	1972-1981	61	303	283	234	112	38	10	0	0	0	0	0	0	0	0	0	0		
	YELLOWTAIL FLOUNDER	1992-2001	918	4,582	4,117	2,767	1,230	230	20	0	0	0	0	0	0	0	0	0	0		
	YELLOWTAIL FLOUNDER	2002-2012	75	374	321	210	65	11	0	0	0	0	0	0	0	0	0	0	0		
	YELLOWTAIL FLOUNDER	SPRING	3,196	15,625	14,140	8,588	3,313	766	133	0	0	0	0	0	0	0	0	0	0		
	YELLOWTAIL FLOUNDER	1963-1971	221	1,062	921	655	314	113	27	0	0	0	0	0	0	0	0	0	0		
	YELLOWTAIL FLOUNDER	1972-1981	530	2,584	2,284	1,671	835	262	64	0	0	0	0	0	0	0	0	0	0		
	YELLOWTAIL FLOUNDER	1982-1991	258	1,240	1,056	680	343	113	22	0	0	0	0	0	0	0	0	0	0		
	YELLOWTAIL FLOUNDER	1992-2001	309	1,524	1,377	832	325	80	13	0	0	0	0	0	0	0	0	0	0		
	YELLOWTAIL FLOUNDER	2002-2012	1,878	9,214	8,502	4,749	1,496	199	7	0	0	0	0	0	0	0	0	0	0		
	YELLOWTAIL FLOUNDER	SUMMER	520	2,529	2,253	1,549	673	166	31	0	0	0	0	0	0	0	0	0	0		
	YELLOWTAIL FLOUNDER	1963-1971	305	1,504	1,360	1,009	428	102	19	0	0	0	0	0	0	0	0	0	0		
	YELLOWTAIL FLOUNDER	1972-1981	200	952	833	523	241	63	12	0	0	0	0	0	0	0	0	0	0		
	YELLOWTAIL FLOUNDER	1982-1991	2	7	6	1	1	0	0	0	0	0	0	0	0	0	0	0	0		
	YELLOWTAIL FLOUNDER	1992-2001	13	66	54	16	3	1	0	0	0	0	0	0	0	0	0	0	0		
	YELLOWTAIL FLOUNDER	FALL	1,588	17,150	15,710	3,359	4,104														

Table 6. Summary of cluster analysis procedures applied to survey catch of juveniles (number) and large spawners (weight).

Procedures run individually on age 0/1 juveniles ¹ and large spawners ²	Process	Sample size or effect	
Hurdle model approach adjustment	Adjust cumulative catch at size, multiplying by the proportion of	All tows included	
Log transform	Transform non-zero catches to a normalized distribution	Zero catches are ignored (reduced number of tows analyzed)	
Select tows for analysis	Select by survey, season, and decade	Reduces number of tows; analysis occurs in desired time period and season; surveys analyzed separately due to catchability differences. Remaining tows may be insufficient number to analyze spatial autocorrelation or hotspots.	
Spatial autocorrelation (Moran's I)	Determine range of highest spatial autocorrelation to set Zone of Indifference parameter for hotspot analysis	Analyzes untransformed tows, including zero catch tows. Procedure may not detect a significant positive spatial autocorrelation. If peak is weak or undetected by analysis, a reasonable alternative was applied for hot spot analysis.	
Hot spot analysis (Getis-Ord's G*) and selection	Identifies hotspots, filtered for significant ($p < 0.05$) hotspots above the mean.	Procedure may not identify any significant hotspots at $p < 0.05$ level.	
Grid hotspots	Number of significant hotspots for a species within a 100 km^2 SASI grid is summed.	All surveys in a season are included, since the hotspot data are standardized relative to each survey's mean.	
Weight layers by importance factor	Number of hotspots in a grid is multiplied by importance factor	Final grid for a season includes all surveys where significant hotspots	

¹ For aged species, upper size threshold that approximated 90th percentile of age 1 fish. Threshold set at the approximate L20 for maturity for unaged species.

² Lower size threshold set where fish at or larger than the threshold comprised 20% of estimated biomass in the spring (applied to spring and summer) and fall (applied to fall and winter) NMFS trawl surveys.

	and summed over species.	were identified by the analysis, weighted by the relative importance of the effect that spatial management will have on regulated groundfish.	
--	--------------------------	---	--

Tows analyzed: Age 0/1 threshold, spring and summer: Age 0/1 threshold, fall and winter: Number of tows with catch:	Tows analyzed: Large spawner threshold: Number of tows with catch:
Spatial autocorrelation	
Freq dist	
Clusters	

		NMFS	MADMF	MENH	IBS cod	IBS yellowtail flounder	IBS Goosefish	
Juvenile	Spring							
	Summer (shrimp or trawl)							
	Summer (dredge)							
	Fall							
	Winter							
Large spawner	Spring							
	Summer (shrimp or trawl)							
	Summer (dredge)							
	Fall							
	Winter							

Table 7. Summary of peak spatial autocorrelation results and alternative trial peaks in parentheses. NA = analysis not attempted due to infrequent catch or data not yet available. NP = No significant peak autocorrelation detected. NSHS = No significant hotspots of above average catches detected or produced by the hotspot analysis. IC = insufficient catch to conduct either a spatial autocorrelation or hotspot analysis.

Species	Survey: NMFS spring		Survey: MADMF spring		Survey: ME/NH spring	
	Juvenile	Spawner	Juvenile	Spawner	Juvenile	Spawner
Cod	8510 (11510)	11510	10528 (15528)	10525 (17528)	4620 (10620)	30620
Haddock	8010 (10010)	8010 (20010)	16528	10528	4620 (6620)	13620 (NSHS)
Yellowtail flounder	11510	11510 (16510)	9528 (14528)	8528 (17528)	IC	IC
American plaice	14510	10510	8528 (17528)	11528	15620	17620
Atlantic wolffish	IC (2 + tows)	NP (20010)	NA	NA	NA	NA
Ocean pout	21510 (12 + tows)	10510	15528 (22528)	13528	5620	17620 NSHS
Pollock	13510	10510	NP (21 + tows)	IC	3620 (7620)	IC
Red Hake	11510 (14510)	NP (14510)	8528	8528	9620	5620
Redfish	9510	10510	IC	11528 (NSHS)	3620 (9620)	4620 (17620) NSHS
Silver hake	10510	32510	20639	10528	6620	11620
White hake	NP (20010)	8510 (21510)	NP (7528)	IC	8620	NP (10620)
Winter flounder	11510	8510 (15510)	7528	8528	3620 (14620)	NP 912620) NSHS
Witch flounder	13510	8510	NP (8528)	IC	7620	NP (3620) NSHS
Windowpane flounder	10510 (23510)	8510	8528 NSHS	8528	4320 NSHS	NP NSHS
Alewife	NA	NA	NA	NA	7620	3620 (20620)
Atlantic herring	NA	NA	NA	NA	4620 (7620)	5620 (23620)
Atlantic halibut	NA	NA	NA	NA	12620	NP NSHS
Goosefish	NA	NA	NA	NA	NA	NA
Barndoor skate	NA	NA	NA	NA	NA	NA

	Survey: IBS Cod spring		Survey: IBS Goosefish spring		Survey: NMFS dredge summer	
Species	Juvenile	Spawner	Juvenile	Spawner	Juvenile	Spawner
Cod	4534 (13534)	NP (28534)	IC	36226	10338 IC	IC
Haddock	11534	7534	NP (48226) NSHS	34226	7338 (16338)	9338 (13338)
Yellowtail flounder	IC	13534 NSHS	IC	34226	5338	5338
American plaice	6534 (9534)	8534	NA	NA	NA	NA
Atlantic wolffish	IC	IC	NA	NA	NA	NA
Ocean pout	IC	IC	NA	NA	NA	NA
Pollock	5334	5334 IC	NA	NA	NA	NA
Red Hake	IC	IC	NA	NA	NP (19338)	IC
Redfish	26534 (5534)	2634 (5534)	NA	NA	NA	NA
Silver hake	IC	IC	NA	NA	NA	NA
White hake	6534 (14534)	6534 (14534)	NA	NA	NA	NA
Winter flounder	5534	5534	NA	NA	16338	17338
Witch flounder	6534 NSHS	6534 NSHS	NA	NA	NA	NA
Windowpane flounder	IC	IC	NA	NA	NA	NA
Alewife	NA	NA	NA	NA	NA	NA
Atlantic herring	NA	NA	NA	NA	NA	NA
Atlantic halibut	NA	NA	NA	NA	NA	NA
Goosefish	NA	NA	35226	NP	NP (19764)	5338 (23338)
Barndoor skate	NA	NA	NA	NA	NP (15338)	11338 (15338)

	Survey: NMFS shrimp summer		Survey: NMFS fall		Survey: MADMF fall	
Species	Juvenile	Spawner	Juvenile	Spawner	Juvenile	Spawner
Cod	8528 (16528)	7528 (13528)	8624 (18624)	8624 (17624)	7365 (9365)	NP (5365) NSHS
Haddock	8528	20528 (26528)	13624	13624	6365 (strong SAC)	22365
Yellowtail flounder	NA	NA	9624	14264	NP (31365) NSHS	4365 (22365) NSHS
American plaice	12528 (18528)	9528 (15528)	9624	10624	4365	6365 (strong peak)
Atlantic Wolffish	IC	IC	IC	NP	IC	IC
Ocean pout	IC	10528 NSHS	24624	9624 (23624)	22365 NSHS	18365 NSHS
Pollock	20528 NSHS	18528 (27528) NSHS	11624 (15624)	8624 (27624)	12365	IC
Red Hake	8528	14528	33624	8624 (33624)	8365	7365 (13365)
Redfish	8528	12528	17624	9624 (17624)	12365 IC	5365
Silver hake	8528 (28528)	9528 (15528)	14624	13624	10365	14365
White hake	12528 (18528)	10528	18624	9624 (18624)	4365	IC
Winter flounder	19528	NC	25624	8624	9365	10365 (22365)
Witch flounder	NP (18528)	8528 (13528)	8964 (22624)	8624 (11624)	17365 IC	10365 NSHS
Windowpane flounder	13528	IC	8964 (22624)	33624	5365 (9365) (strong 2 nd peak)	6365 (10365) (strong 2 nd peak)
Alewife	NA	NA	NA	NA	NA	NA
Atlantic herring	NP	NP	12624	16624	11365	5365 NSHS
Atlantic halibut	NA	NA	NA	NA	NA	NA
Goosefish	11528	28528 NSHS	13624	9624 (12624) NSHS	11365 (13365) NSHS	5365 NSHS
Barndoor skate	NA	NA	NA	NA	NA	NA

	Survey: ME/NH fall		Survey: IBS Cod fall		Survey: IBS YTF fall	
Species	Juvenile	Spawner	Juvenile	Spawner	Juvenile	Spawner
Cod	5988 (7988)	4988 (21998)	7313	9313	IC	IC
Haddock	29998	NP IC	7313	20913	IC	IC
Yellowtail flounder	8988 NSHS	NP IC	IC	5313	24642 NSHS	16642
American plaice	24988	3988	5313	NP (25313)	NA	NA
Atlantic wolffish	NA	NA	IC	IC	NA	NA
Ocean pout	4998	IC	NA	NA	NA	NA
Pollock	NP (18998)	IC	NP (11313) NSHS	12313	NA	NA
Red Hake	16998 (strong peak)	10998 (strong peak)	IC	IC	NA	NA
Redfish	5998 (17998)	NP 6998	12313	NP (8313)	NA	NA
Silver hake	13998	9988	IC	IC	NA	NA
White hake	17998	6998 IC	10313	IC	NA	NA
Winter flounder	17998	NP IC	5313 (17313)	7313	IC	IC
Witch flounder	4998 (14998)	8998 (17998) NSHS	NP	5313 (9313)	NA	NA
Windowpane flounder	8988	3988 IC	IC	7313	NA	NA
Alewife	16988	7988 (17988)	NA	NA	NA	NA
Atlantic herring	5998	3988	NA	NA	NA	NA
Atlantic halibut	12998 IC	3998 IC	NA	NA	NA	NA
Goosefish	11998 NSHS	IC	5313 (9313)	NP (23313)	NP IC	IC
Barndoor skate	NA	NA	NA	NA	NA	NA


	Survey: NMFS winter		Survey: IBS Cod winter		Survey: IBS GSF winter	
Species	Juvenile	Spawner	Juvenile	Spawner	Juvenile	Spawner
Cod	15806	27806	9728 (12728)	NP (7728)	NP (31083) NSHS	NP
Haddock	17806	NP (23806)	17728 (31728)	10728	NP	49083
Yellowtail flounder	21806	12806 (28806)	IC	NP (3728)	IC	NP
American plaice	IC	24806	8728	6728	59083 NSHS	35083 NSHS
Atlantic wolffish	NA	NA	IC	IC	NA	NA
Ocean pout	14806 (16806)	14806	IC	IC	NA	NA
Pollock	IC	IC	IC	NP (15728)	NA	NA
Red Hake	20806 (27806)	12806	NA	NA	NA	NA
Redfish	NA	NA	NA	NA	NA	NA
Silver hake	19806	12806 (31806)	NA	NA	NA	NA
White hake	NA	NA	11728	NP IC		
Winter flounder	12806 (16806)	21806	5728 (20728)	NP (24728) NSHS	35083	NP NSHS
Witch flounder	19806	12806 (14806)	7728 (12728)	8728	IC	36083 (40083)
Windowpane flounder	15806 (17806)	14806 (37806)	IC	6728	NA	NA
Alewife	NA	NA	NA	NA	NA	NA
Atlantic herring	NA	NA	NA	NA	NA	NA
Atlantic halibut	NA	NA	NA	NA	NA	NA
Goosefish	12806 (25806)	32806	6728 (21728)	NP	35083 (44083)	34083
Barndoor skate	NA	NA	NA	NA	40083 NSHS	NP NSHS

Table 8. Summary of significant hotspots of above average catches identified by survey and species for age 0/1 juvenile (upper) and for large spawners (lower), 2002-2012.

Survey	Years	Tows	Mean to ne:	StdDev			Alewife	Am Pleice	Atlantic Herring	Atlantic halibut	Atlantic Wolffish	Barndoor skate	Cod	Goosefish	Haddock	Ocean pout	Pollock	Red hake	Redfish	Silver hake	White hake	Winter flounder	Witch flounder	Windowpane flounder	Yellowtail flounder	Total survey	
					90th pctle	95th pctle																					
NMFS spring	2002-2012	3,426	4,012.0	3,630.0	7,509.5	9,014.9		85	0				35	31	0	0	122	25	167	70	53	7	3	11	609		
NMFS shrimp		677	3,088.9	2,328.5	6,527.5	8,258.9		114					1	48	4		23	161	87	112		56		7	431		
NMFS scallop	2002-2011	4,634	1,538.7	1,454.9	3,337.7	4,269.8						81	18	250	61			0				14			5	1082	
NMFS fall	2002-2011	3,413	4,004.0	2,634.0	7,624.0	8,979.0		91	1				33	30	80	0	1	286	69	254	77	132	19	4	5	99	
NMFS winter	2002-2007	659	6,212.4	5,272.9	11,805.6	13,468.3		0					2	3	1	1		18		59		8	3	4	0	17	
MADMF spring	2002-2012	936	832.9	655.3	1,798.9	2,184.9							80		8	0	3	19	0	41	4	150		0	17	366	
MADMF fall	2002-2011	714	1,096.8	835.9	2,364.8	2,807.9							5	0	4	0	0	58	0	88	2	131		2	315		
MENH spring		1,194	1,078.7	1,156.7	2,619.4	3,298.2	187	269	51	19			85		36	9	16	70	116	317	71	264	57	149	0	1716	
MENH fall		812	1,271.7	1,436.0	2,987.9	3,859.1	192	233	92	11			29	0	15	4	4	186	329	275	209	187	46	134	0	1946	
IBS cod spring		449	1,513.1	1,643.0	3,533.9	4,638.3			77				54		25				18		10	16	0		200		
IBS cod fall		175	2,202.4	2,559.9	4,312.8	6,101.3			12				21	7	8		0		2		8	28	0		86		
IBS cod winter		274	2,064.9	3,114.4	3,728.0	5,131.3							2	10	10					14	65	1		102			
IBS goosefish spring		229	15,551.0	13,125.6	30,226.1	34,028.5								13	0										13		
IBS goosefish winter		198	16,992.9	9,778.9	31,082.6	34,286.3							2	0											2		
IBS YTF fall		709	3,382.5	14,471.1	5,642.0	7,373.3																		0	0		
Total species hotspots =								379	949	145	30	0	81	367	361	283	14	24	782	720	1288	577	1048	189	296	40	7573

Survey	Years	Tows	Mean to ne:	StdDev			Alewife	Am Pleice	Atlantic Herring	Atlantic halibut	Atlantic Wolffish	Barndoor skate	Cod	Goosefish	Haddock	Ocean pout	Pollock	Red hake	Redfish	Silver hake	White hake	Winter flounder	Witch flounder	Windowpane flounder	Yellowtail flounder	Total survey	
					90th pctle	95th pctle																					
NMFS spring	2002-2012	3,426	4,012.0	3,630.0	7,509.5	9,014.9		43	67				22	145	14	6	92	19	174	7	5	4	35	30	663		
NMFS shrimp		677	3,088.9	2,328.5	6,527.5	8,258.9		23	66				1	0	16	0	0	139	71		0	4			320		
NMFS scallop	2002-2011	4,634	1,538.7	1,454.9	3,337.7	4,269.8						1	1		3							24		17	46		
NMFS fall	2002-2011	3,413	4,004.0	2,634.0	7,624.0	8,979.0		14					16	0	91	1	13	259	51	141	13		4	51	39	693	
NMFS winter	2002-2007	659	6,212.4	5,272.9	11,805.6	13,468.3							0	3	1	14		2		31			0	20	3	74	
MADMF spring	2002-2012	936	832.9	655.3	1,798.9	2,184.9	127	3					0		1	30		9		24		5		29	29	257	
MADMF fall	2002-2011	714	1,096.8	835.9	2,364.8	2,807.9	1	0					0	0	0	0	0	24		30		0	2		57		
MENH spring		1,194	1,078.7	1,156.7	2,619.4	3,298.2	73	74	0	0			0		0	0		15	0	38		0	0		0	200	
MENH fall		812	1,271.7	1,436.0	2,987.9	3,859.1	19	2	23	0			2					57	39	54	0		0	0	0	196	
IBS cod spring		449	1,513.1	1,643.0	3,533.9	4,638.3			14				7		28		1		6			0	0	1	0	57	
IBS cod fall		175	2,202.4	2,559.9	4,312.8	6,101.3			0				8	0	6		0		1				3	2	4	24	
IBS cod winter		274	2,064.9	3,114.4	3,728.0	5,131.3			2				6	0	13		4					0	0	3	28		
IBS goosefish spring		229	15,551.0	13,125.6	30,226.1	34,028.5							1	1	5										7		
IBS goosefish winter		198	16,992.9	9,778.9	31,082.6	34,286.3			5				0	0	2	3						4	0	0	14		
IBS YTF fall		709	3,382.5	14,471.1	5,642.0	7,373.3							0									65		65			
Total species hotspots =								146	180	230	0	0	1	64	6	312	59	24	597	187	492	20	38	15	140	190	2701


Figure 2. Data processing flowchart for spatial autocorrelation and hotspot analyses for juvenile (upper) and large spawner (lower) life stages. The example analyzes witch flounder juvenile and large spawner distribution in the 2009 IBS winter goosefish survey.



Summary of hotspot analysis parameters and results
April 4, 2013

Figure 3. Workflow for merging and gridding weighted number of hotspots for a season.

Merge survey layers and grid for a species	Apply grid weights and merge for a season
--	---



Getis-Ord Gi* statistic

The [Hot Spot Analysis](#) tool calculates the Getis-Ord Gi* statistic (pronounced G-i-star) for each feature in a dataset. The resultant [z-scores and p-values](#) tell you where features with either high or low values cluster spatially. This tool works by looking at each feature within the context of neighboring features. A feature with a high value is interesting but may not be a statistically significant hot spot. To be a statistically significant hot spot, a feature will have a high value and be surrounded by other features with high values as well. The local sum for a feature and its neighbors is compared proportionally to the sum of all features; when the local sum is very different from the expected local sum, and that difference is too large to be the result of random chance, a statistically significant [z-score](#) results.

Calculations

The Getis-Ord local statistic is given as:

$$G_i^* = \frac{\sum_{j=1}^n w_{i,j} x_j - \bar{X} \sum_{j=1}^n w_{i,j}}{S \sqrt{\left[n \sum_{j=1}^n w_{i,j}^2 - \left(\sum_{j=1}^n w_{i,j} \right)^2 \right] / (n-1)}} \quad (1)$$

where x_j is the attribute value for feature j , $w_{i,j}$ is the spatial weight between feature i and j , n is equal to the total number of features and:

$$\bar{X} = \frac{\sum_{j=1}^n x_j}{n} \quad (2)$$

$$S = \sqrt{\frac{\sum_{j=1}^n x_j^2}{n} - (\bar{X})^2} \quad (3)$$

The G_i^* statistic is a z-score so no further calculations are required.

Interpretation

The Gi* statistic returned for each feature in the dataset is a z-score. For statistically significant positive z-scores, the larger the z-score is, the more intense the clustering of high values (hot spot). For statistically significant negative z-scores, the smaller the z-score is, the more intense the clustering of low values (cold spot). For more information about determining statistical significance, see [What is a z-score? What is a p-value?](#)

Output

This tool creates a new **Output Feature Class** with a z-score and p-value for each feature in the **Input Feature Class**. If there is a selection set applied to the Input Feature Class, only selected features will be analyzed, and only selected features will appear in the Output Feature Class. This tool also returns the z-score and p-value field names as derived output values for potential use in custom models and scripts.

When this tool runs in ArcMap, the **Output Feature Class** is automatically added to the table of contents with default rendering applied to the z-score field. The hot to cold rendering applied is defined by a layer file in <ArcGIS>/ArcToolbox/Templates/Layers. You can reapply the default rendering, if needed, by [importing](#) the template layer symbology.


Hot spot analysis considerations

There are three things to consider when undertaking any hot spot analysis:

1. What is the Analysis Field (**Input Field**)? The hot spot analysis tool assesses whether high or low values (the number of crimes, accident severity, or dollars spent on sporting goods, for example) cluster spatially. The field containing those values is your Analysis Field. For point incident data, however, you may be more interested in assessing incident intensity than in analyzing the spatial clustering of any particular value associated with the incidents. In that case, you will need to aggregate your incident data prior to analysis. There are several ways to do this:
 - If you have polygon features for your study area, you can use the [Spatial Join](#) tool to count the number of events in each polygon. The resultant field containing the number of events in each polygon becomes the **Input Field** for analysis.
 - Use the [Create Fishnet](#) tool to construct a polygon grid over your point features. Then use the [Spatial Join](#) tool to count the number of events falling within each grid polygon. Remove any grid polygons that fall outside your study area. Also, in cases where many of the grid polygons within the study area contain zeros for the number of events, increase the polygon grid size, if appropriate, or remove those zero-count grid polygons prior to analysis.
 - Alternatively, if you have a number of coincident points or points within a short distance of one another, you can use [Integrate](#) with the [Collect Events](#) tool to (1) snap features within a specified distance of each other together, then (2) create a new feature class containing a point at each unique location with an associated count attribute to indicate the number of events/snapped points. Use the resultant ICOUNT field as your **Input Field** for analysis.

Note:

If you are concerned that your coincident points may be redundant records, the [Find Identical](#) tool can help you to locate and remove duplicates.



Strategies for aggregating incident data

2. Which **Conceptualization of Spatial Relationships** is appropriate? What **Distance Band or Threshold Distance** value is best?

The recommended (and default) **Conceptualization of Spatial Relationships** for the [Hot Spot Analysis \(Getis-Ord Gi*\)](#) tool is **Fixed Distance Band**. Space-Time Window, Zone of Indifference, Contiguity, K Nearest Neighbor, and Delaunay Triangulation may also work well. For a discussion of best practices and strategies for determining an analysis distance value, see [Selecting a Conceptualization of Spatial Relationships](#) and [Selecting a Fixed Distance](#). For more information about space-time hot spot analysis, see [Space-Time Analysis](#).

3. What is the question?

This may seem obvious, but how you construct the **Input Field** for analysis determines the types of questions you can ask. Are you most interested in determining where you have lots of incidents, or where high/low values for a particular attribute cluster spatially? If so, run [Hot Spot Analysis](#) on the raw values or raw incident counts. This type of analysis is particularly helpful for resource allocation types of problems. Alternatively (or in addition), you may be interested in locating areas with unexpectedly high values in relation to some other variable. If you are analyzing foreclosures, for example, you probably expect more foreclosures in locations with more homes (said another way, at some level, you expect the number of foreclosures to be a function of the number of houses). If you divide the number of foreclosures by the number of homes, then run the Hot Spot Analysis tool on this ratio, you are no longer asking Where are there lots of foreclosures?; instead, you are asking Where are there unexpectedly high numbers of foreclosures, given the number of homes? By creating a rate or ratio prior to analysis, you can control for certain expected relationships (for example, the number of crimes is a function of population; the number of foreclosures is a function of housing stock) and identify unexpected hot/cold spots.

Best practice guidelines

- Does the **Input Feature Class** contain at least 30 features? Results aren't reliable with less than 30 features.
- Is the **Conceptualization of Spatial Relationships** you selected appropriate? For this tool, the **Fixed Distance Band** method is recommended. For space-time hot spot analysis, see [Selecting a Conceptualization of Spatial Relationships](#).
- Is the **Distance Band or Threshold Distance** appropriate? See [Selecting a Fixed Distance](#).
 - All features should have at least one neighbor.
 - No feature should have all other features as neighbors.
 - Especially if the values for the **Input Field** are skewed, you want features to have about eight neighbors each.

Potential applications

Applications can be found in crime analysis, epidemiology, voting pattern analysis, economic geography, retail analysis, traffic incident analysis, and demographics. Some examples include the following:

- Where is the disease outbreak concentrated?
- Where are kitchen fires a larger than expected proportion of all residential fires?
- Where should the evacuation sites be located?
- Where/When do peak intensities occur?
- Which locations and at during what time periods should we allocate more of our resources?

Additional resources

Mitchell, Andy. *The ESRI Guide to GIS Analysis*, Volume 2. ESRI Press, 2005.

Getis, A. and J.K. Ord. 1992. "The Analysis of Spatial Association by Use of Distance Statistics" in *Geographical Analysis* 24(3).

Ord, J.K. and A. Getis. 1995. "Local Spatial Autocorrelation Statistics: Distributional Issues and an Application" in *Geographical Analysis* 27(4).

How Incremental Spatial Autocorrelation works


[Desktop](#) » [Geoprocessing](#) » [Tool reference](#) » [Spatial Statistics toolbox](#) » [Analyzing Patterns toolset](#)

With much of the spatial data analysis you do, the scale of your analysis will be important. The default **Conceptualization of Spatial Relationships** for the [Hot Spot Analysis](#) tool, for example, is **FIXED_DISTANCE_BAND** and requires you to specify a distance value. For many density tools you will be asked to provide a **Radius**. The distance you select should relate to the scale of the question you are trying to answer or to the scale of remediation you are considering. Suppose, for example, you want to understand childhood obesity. What is your scale of analysis? Is it at the individual household or neighborhood level? If so, the distance you use to define your scale of analysis will be small, encompassing the homes within a block or two of each other. Alternatively, what will be the scale of remediation? Perhaps your question involves where to increase after-school fitness programs as a way to potentially reduce childhood obesity. In that case, your distance will likely be reflective of school zones. Sometimes it's fairly easy to determine an appropriate scale of analysis; if you are analyzing commuting patterns and know that the average journey to work is 12 miles, for example, then 12 miles would be an appropriate distance to use for your analysis. Other times it is more difficult to justify any particular analysis distance. This is when the [Incremental Spatial Autocorrelation](#) tool is most helpful.

Whenever you see spatial clustering in the landscape, you are seeing evidence of underlying spatial processes at work. Knowing something about the spatial scale at which those underlying processes operate can help you select an appropriate analysis distance. The [Incremental Spatial Autocorrelation](#) tool runs the [Spatial Autocorrelation \(Global Moran's I\)](#) tool for a series of increasing distances, measuring the intensity of spatial clustering for each distance. The intensity of clustering is determined by the [z-score](#) returned. Typically, as the distance increases, so does the z-score, indicating intensification of clustering. At some particular distance, however, the z-score generally peaks.

Sometimes you will see multiple peaks.

Spatial Autocorrelation by Distance



Peaks reflect distances where the spatial processes promoting clustering are most pronounced. The color of each point on the graph corresponds to the statistical significance of the [z-score](#) values.

Significance Level (p-value)	Critical Value (z-score)
0.01	< -2.58
0.05	-2.58 - -1.96
0.10	-1.96 - -1.65
---	-1.65 - 1.65
0.10	1.65 - 1.96
0.05	1.96 - 2.58
0.01	> 2.58

One strategy for identifying an appropriate scale of analysis is to select the distance associated with the [statistically significant](#) peak that best reflects the scale of your question. Often this is the first statistically significant peak.

How do I select the Beginning Distance and Distance Increment values?

All distance measurements are based on feature centroids and the default **Beginning Distance** is the smallest distance that will ensure every feature has at least one neighboring feature. This is generally a good choice, unless your dataset includes spatial outliers. [Determine whether or not you have spatial outliers](#), then select all but the outlier features and [run Incremental Spatial Autocorrelation on just the selected features](#). If you find a peak distance for the selection set, [use that distance](#) to create a [spatial weights matrix file](#) based on all of your features (even the outliers). When you run the [Generate Spatial Weights Matrix](#) tool to create the spatial weights matrix file, set the **Number of Neighbors** parameter to some value so that all features will have [at least that many neighboring features](#).

The default **Increment Distance** is the average distance to each feature's nearest neighboring feature. If you've determined an appropriate starting distance using the strategies above and still don't see a peak distance, you may want to experiment with smaller or larger increment distances.

What if the graph never peaks?


In some cases, you will use the [Incremental Spatial Autocorrelation](#) tool and get a graph with a [z-score](#) that just continues to rise with increasing distances; there is no peak. This most often happens in cases where data has been aggregated and the scale of the processes impacting your **Input Field** variable are smaller than the aggregation scheme. You can try making your Distance Increment smaller to see if this captures more subtle peaks. Sometimes, however, you won't get a peak because there are multiple spatial processes, each operating at a different distance, in your study area. This is often the case with large point datasets that are noisy (no clear spatial pattern to the point data values you're analyzing). In this case, you will need to justify your scale of analysis using some other criteria.

Interpreting results

When you run the [Incremental Spatial Autocorrelation](#) tool in the [foreground](#), the z-score results for each distance are written to the [Progress](#) window. This output is also available from the [Results window](#). If you right-click on the [Messages entry](#) in the [Results window](#) and select [View](#), the tool results are displayed in a [Message](#) dialog box. When you specify a path for the optional [Output Table](#) parameter, a table is created that includes fields for **Distance**, **MoransI**, **ExpectedI**, **Variance**, **z_score**, and **p_value**. By examining the z-score values in the [Progress](#) window, [Message](#) dialog box, or [Output Table](#), you can determine if there are any peak distances. More typically, however, you would identify peak distances by looking at the graphic in the optional [Output Report](#) file. The report has three pages. An example of the first page of

the report is shown below. Notice that this graph has three peak z-scores associated with distances of 5000, 9000, and 13000 feet. A halo will be drawn to highlight both the first peak distance and the maximum peak distance, but all peaks represent distances where the spatial processes promoting clustering are most pronounced. You can select the peak that best reflects the scale of your analytical question. In some cases, there will only be one halo because the first and the maximum peaks are found at the same distance. If none of the z-score peaks are statistically significant, then none of the peaks will have the light blue halo. Notice that the color of the plotted z-score corresponds to the legend showing the critical values for statistical significance.

Spatial Autocorrelation by Distance



On page two of the report, the distances and z-score values are presented in table format. The last page of the report documents the parameter settings used when the tool was run. To get a report file, provide a path for the **Output Report** parameter.

Figure 4. Example of ‘good’ spatial autocorrelation result: Large spawner silver hake from MADMF fall survey, 2002-2011.




Figure 5. Example of ‘satisfactory’ spatial autocorrelation result, with secondary peak autocorrelation: Juvenile American plaice from IBS cod fall survey, 2002-2011.




Figure 6. Example of unsatisfactory spatial autocorrelation result, with no significant peak in autocorrelation: Large spawner American plaice from IBS cod fall survey, 2002-2011. In this case, hotspot analysis was re-run with a zone of indifference parameter of 25313 m, corresponding of a secondary non-significant spatial autocorrelation peak, but there were no significant hotspots identified nonetheless.




Figure 7. Example of unsatisfactory spatial autocorrelation resulting from insufficient non-zero catches: Large spawner pollock from IBS cod fall survey, 2002-2011. No significant hotspots were identified and no further analysis was attempted.




Figure 8. Example of ‘good’ spatial autocorrelation result, but first autocorrelation peak is probably not meaningful: Juvenile winter flounder from IBS cod fall survey, 2002-2011. The maximum peak of 17,313 m was used as the Zone of Indifference parameter in the hotspot analysis in lieu of the first peak.




Figure 9. Example of unsatisfactory spatial autocorrelation: Juvenile witch flounder from IBS cod fall survey, 2002-2011. No significant hotspots were identified and no further analysis was attempted.




Figure 10. Example of ‘good’ spatial autocorrelation result, with no meaningful first autocorrelation: Large spawner yellowtail flounder from NMFS winter survey, 2002-2007. The maximum peak was applied as a Zone of Indifference parameter in the hotspot analysis.




Figure 11. Example of ‘poor’ spatial autocorrelation result. Data are sparse and tend the spatial autocorrelation has a ‘choppy’ appearance: Juvenile cod from NMFS winter survey, 2002-2007. Usually, this pattern is associated with a hotspot analysis that has no significant positive hotspots.




Figure 12. Example of ‘strong’ spatial autocorrelation result: Large spawner witch flounder from the NMFS winter survey, 2002-2007.

